

MATTEO BONFORTE and AGNESE DI CASTRO

Quantitative local estimates for nonlinear elliptic equations involving p -Laplacian type operators

Abstract. The purpose of this paper is to prove quantitative local upper and lower bounds for weak solutions of elliptic equations of the form $-\Delta_p u = \lambda u^s$, with $p > 1$, $s \geq 0$ and $\lambda \geq 0$, defined on bounded domains of \mathbb{R}^d , $d \geq 1$, without reference to the boundary behaviour. We give an explicit expression for all the involved constants. As a consequence, we obtain local Harnack inequalities with explicit constants. Finally, we discuss the issue of local absolute bounds, which are new to our knowledge. Such bounds will be true only in a restricted range of s or for a special class of weak solutions, namely for local stable solutions. In the study of local absolute bounds for stable solutions there appears the so-called Joseph-Lundgren exponent as a limit of applicability of such bounds.

Keywords. Nonlinear elliptic equations of p -Laplacian type, local bounds, Harnack inequalities.

Mathematics Subject Classification (2010): 35B45, 35B65, 35J60, 35J61.

Contents

1 - Introduction	214
1.1 - Plan of the paper and main results	216
1.2 - More general nonlinearities	217
2 - Local energy estimates and Caccioppoli inequalities	218
3 - Sobolev and reverse Poincaré inequalities	222

4 - Local upper bounds	227
4.1 - <i>Local upper bounds I. The upper Moser iteration</i>	228
4.2 - <i>Local upper bounds II. The case of unbounded coefficients</i>	238
5 - Local lower bounds via Moser iteration	244
5.1 - <i>Reverse Hölder inequalities and additional local lower bounds</i>	248
6 - Harnack inequalities	251
7 - Local absolute bounds	257
7.1 - <i>Local absolute bounds for $s < s_c^*$</i>	257
7.2 - <i>Local absolute bounds for stable solutions. The supercritical case ..</i>	258
8 - Appendix	264
8.1 - <i>The John-Nirenberg Lemma and reverse Hölder inequalities</i>	264
8.2 - <i>Technical tools</i>	266
8.3 - <i>Numerical identities and inequalities</i>	267

1 - Introduction

In this paper we obtain local upper and lower estimates for the weak solutions of nonlinear elliptic equations of the form

$$(1) \quad -\Delta_p u = -\operatorname{div}(|\nabla u|^{p-2} \nabla u) = f(u),$$

with $p > 1$, posed in a bounded domain $\Omega \subset \mathbb{R}^d$, with $d \geq 1$. The choice of right-hand side that we have in mind is $f(u) = \lambda u^s$ with $\lambda, s > 0$. Our main purpose is to obtain local estimates for solutions that are defined inside the domain without reference to their boundary behaviour. The notion of solution that we will use in the whole paper is the following.

Definition 1.1 (Local weak solutions). Let $\Omega \subset \mathbb{R}^d$ be a bounded domain. A function u is a *local weak solution* to $-\Delta_p u = f(u)$ in Ω if and only if $u \in W_{\text{loc}}^{1,p}(\Omega)$, $f(u) \in L_{\text{loc}}^1(\Omega)$ and it satisfies

$$\int_{\Omega} \left[|\nabla u|^{p-2} \nabla u \cdot \nabla \varphi - f(u) \varphi \right] dx = 0, \quad \text{for all } \varphi \in C_c^1(\Omega).$$

Throughout the paper, $C_c^1(\Omega)$ will be the space of $C^1(\Omega)$ functions with compact support in Ω .

The estimates that we prove in this paper are local upper bounds for solutions of any sign, lower bounds for nonnegative solutions, and also local Harnack inequalities. The estimates that we obtain are not essentially new from a qualitative point of view, and enjoy a large literature [2, 4, 5, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 28, 29, 31, 32, 37, 39, 41, 47, 48, 51, 52, 53, 54, 55] and the books [33, 34, 36, 49]; however, it

is hopeless to give a complete bibliography for this nowadays classical problem. We try to contribute to the general theory on these elliptic equations, with quantitative local bounds; to our knowledge, there does not exist in literature a systematic set of local quantitative estimates in the explicit form given here. By quantitative estimates we mean keeping track of all the constants during the proofs. This paper follows the ideas of [4], in which the authors treated the case $p = 2$, i. e. the case of semilinear equations. Here we extend the techniques and the results of [4] to the more general case represented by the p -Laplacian elliptic equation (1).

The interest in obtaining quantitative control of the constants of such inequalities relies in the applications. On one hand, our results are useful in understanding regularity properties of the stationary solutions of the associated parabolic equation (the so-called doubly nonlinear evolution equation); it is needed for instance in the results of [3] on the asymptotic properties of solutions of the fast diffusion equation in bounded domains. On the other hand, it is interesting to see the stability of the estimates (therefore of the regularity of the solutions), when the parameters s or p reach their limiting values; for example, we can consider the (formal) limit $p \rightarrow 1^+$ in the local upper estimates of Theorems 4.1 and 4.4 and easily check that the constant is stable under such limiting process; the upper estimates therefore should hold also for the solutions of the 1-Laplacian, often called the Total Variation Flow (TVF). Weak solutions to the TVF have a different definition from the one we provide here for the p -Laplacian, but are sometimes obtained as the limit for $p \rightarrow 1^+$ of suitable families $\{u_p\}$ of smooth solution to the p -Laplacian, see [1] for more details; we refrain from doing this limiting process, since it falls out from the scope of this paper.

The range of exponents of interest will be $p > 1$ and $0 \leq s < r - 1$, where r is the exponent of the Sobolev imbedding of $W^{1,p}$, namely $r = p^* = pd/(d - p)$ if $p < d$ and any $r \in [p^*, \infty)$ if $p \geq d$; it is clear at this point that there is a restriction on the parameter s only when $p < d$. It is worth noticing that the restriction $s < r - 1 = p^* - 1$ appears only when we consider $p < d$, and it is related to several deep aspects of the theory of the equation at hand: for example, when dealing with the homogeneous Dirichlet problem, the existence of bounded weak solutions may fail above that exponent, as well as the absolute upper bounds, see [11, 23, 31, 32, 46, 52]; it is known that when $s \geq p^* - 1$ there exist solutions¹ which are not bounded, therefore not regular, cf. [24, 40, 42, 43, 44, 45]. On the other hand, when $s < p^* - 1$, bounded solutions are known to be $C^{1,\alpha}$, cf. [25], and the $C^{1,\alpha}$ modulus of continuity directly depends on the local maximum of the solution. Therefore having absolute bounds for the solution allows to have

¹ for $p = 2$, very weak solutions.

absolute bounds for the $C^{1,\alpha}$ modulus of continuity. We will see that the $C^{1,\alpha}$ modulus of continuity is independent of the solution when $s < s_c^* < p^* - 1$, while it depends on (some L^q -norms of) the solution when $s_c^* < s < p^* - 1$. Finally, when dealing with quantitative local absolute bounds for the smaller class of stable solutions, cf. Section 7, there will naturally appear the so-called Joseph-Lundgren exponent s_{JL} – which is finite only for “large” dimensions – as a further limit on the range of s to which our absolute bounds apply, as we shall explain in Section 7. As a reference for this topic see for example [13, 14, 16, 28, 29].

1.1 - Plan of the paper and main results

We begin with a section devoted to the basic energy estimates. As a consequence, we obtain quantitative Caccioppoli type estimates that allow us to obtain absolute bounds for the $s - 1$ -“norm”, which to our knowledge have never been observed before, see Corollary 2.1 ; such absolute bounds will be the key tool needed in Section 7 to derive our local absolute bounds. In Section 3 we recall the Sobolev inequalities that we will use in the paper and derive some preliminary inequalities in the form of reverse Poincaré inequalities, as a direct consequence of the energy estimates and Sobolev inequalities. We then focus on local upper estimates in Section 4. Our first main result is Theorem 4.1, which can be considered as a smoothing effect with very precise constants. In the case $p - 1 < s < r - 1$, the estimates of Theorem 4.1 seem to be new to our knowledge. Next, we obtain local upper estimates for $-\Delta_p u = b u^{p-1}$ with unbounded coefficient b in Theorems 4.2 and 4.3 and we apply them to the case $b = u^{s-(p-1)}$ in Theorem 4.4. The last upper bounds have the advantage that they do not require the restriction $s < r - 1$, they hold for any nonnegative weak solution which moreover belongs to L_{loc}^q , with $q > r[s - (p - 1)]/(r - p)$. This last requirement seems to be essential, since in the case $s > r - 1$ there are solutions u_∞ which are not bounded, and $u_\infty \in L_{loc}^q$ with $q < r[s - (p - 1)]/(r - p)$, at least when $p = 2$, see [24, 40, 42, 43, 44, 45].

Section 5 is devoted to the local lower estimates. The main result is Theorem 5.1, which holds for all $p > 1$ and $0 \leq s < r - 1$. The proof is based on a quantitative lower Moser iteration, joined with the reverse Hölder inequalities of Appendix 8.1, which are obtained via a simplified John-Nirenberg type Lemma proved in [4] in a quantitative form. Next we prove a more precise quantitative reverse Hölder inequality, Proposition 5.1, but only in the smaller range of exponents $p - 1 < s < r(p - 1)/p = s_c^*$. The fourth main result of the paper, is Theorem 5.2, in which we use such reverse Hölder inequality to improve the lower bounds of Theorem 5.1 in this smaller range of exponents.

In Section 6 we combine the upper bounds of Section 4 with the lower bounds of Section 5 to obtain various form of Harnack inequalities. The general form, valid in the whole range of exponents, is given in Theorem 6.1, but, unfortunately, the constant of such inequality, depends on a quotient of L^q norms. Next we specialize to the subcritical range $0 < s \leq p - 1$, Theorem 6.2, and supercritical range $p - 1 < s < s_c^*$, Theorem 6.3, and we prove clean versions of the Harnack inequality, i.e. the constant is independent of the solution. In the range $s_c^* < s < r - 1$, we are not able to prove such clean forms of Harnack inequalities, and we conjecture that the dependence on some L^q norm of the solution can not be avoided. As far as we know, the Harnack inequality that we derive for $s > p - 1$ is not stated explicitly in the literature. The fact that the “constant” involved has to depend on u when $s_c^* \leq s < r - 1$ is confirmed by the results of [9, 6, 7, 8, 26, 27] applied to separation of variable solutions of parabolic problems. This is also related to the fact that, in the range $s_c^* \leq s < r - 1$, there exist (very weak) singular solutions, at least when $p = 2$, see [24, 40, 42, 43, 44, 45].

Finally, in Section 7 we derive the quantitative local absolute bounds, which represent the novelty of the paper. In Theorem 7.1 we obtain quantitative local lower bounds when $0 \leq s < p - 1$ and local absolute upper bounds when $p - 1 < s < s_c^*$. We have already discussed why the above absolute bounds cannot be extended $s > s_c^*$ without further assumption on the solution. The last part of the section is devoted to the derivation of absolute upper bounds for all $s \geq 0$, but for the class of local stable solutions. In Theorem 7.2 we obtain quantitative absolute upper bounds for all $s > 0$ when the dimension is small, namely $d \leq \frac{p(p+3)}{p-1}$, while we reach a bigger exponent $s_{JL} \in (r - 1, \infty)$ for bigger dimension. The exponent s_{JL} is the celebrated exponent discovered by Joseph and Lundgren in [35], see also in [13, 14, 16, 28, 29]. The Appendix contains some technical results used in the paper, complemented with a proof when needed. We will use the notation $\|g\|_{L^q(B_\rho)} = \|g\|_{q,\rho}$, $|B_\rho| = \omega_d \rho^d$ and $\omega_d = |B_1|$.

1.2 - More general nonlinearities

We can apply the method used in the proofs to obtain quantitative estimates to a larger class of operators and nonlinearities. We can consider a more general equation, namely

$$A(u) = -\operatorname{div} a(x, u, \nabla u) = f(u),$$

where $a(x, \sigma, \zeta)$ is a Carathéodory vector valued function on $\Omega \times \mathbb{R} \times \mathbb{R}^d$ such that, for some constants $v_1 \geq v_2 > 0$

1. $|a(x, \sigma, \xi)| \leq v_1[1 + |\xi|^{p-1}]$,
2. $a(x, \sigma, \xi)\xi \geq v_2|\xi|^p$,
3. $[a(x, \sigma, \xi) - a(x, \sigma, \eta)][\xi - \eta] > 0$,

for a. e. $x \in \Omega$ and $\forall \sigma \in \mathbb{R}, \xi, \eta \in \mathbb{R}^d, \xi \neq \eta$.

The proofs of all the results apply also to this case with minor modifications, but the constants in the estimates will also depend on v_1 and v_2 . As far as the the right-hand side is concerned, we deal with the model case $f(u) = \lambda u^s$. Indeed, we could have considered a more general nonlinearity $f(u)$ satisfying the following conditions: there exist $0 \leq b_0 \leq b_1, b_2 \geq 0$:

$$b_0 u^s \leq f(u) \leq b_1(u + b_2)^s.$$

Also in this case the proofs of all the results apply with minor modifications, and it is not so difficult to keep track of the new constants b_i throughout the proof.

We have decided here to consider the model case, to simplify the exposition and to focus on the main ideas.

2 - Local energy estimates and Caccioppoli inequalities

We shall pursue in the sequel the well-known idea that local weak solutions satisfy reverse Sobolev or Poincaré inequalities. Such local reverse inequalities are the key to prove local upper and lower estimates of next sections, and indeed imply that such functions satisfy Harnack inequalities.

Lemma 2.1 (Energy Estimates). *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain and let u be a local nonnegative weak solution to $-\Delta_p u = \lambda u^s$ in Ω , $p > 1$ and $\lambda, s \geq 0$. Then the following energy estimate holds true for any $\alpha \neq -(p - 1)$, $\delta > 0$ and any test function $\phi \in C_c^1(\Omega)$, $\phi > 0$*

$$\begin{aligned} (2) \quad & \frac{|\alpha|}{p} \left| \frac{p}{\alpha + (p - 1)} \right|^p \int_{\Omega} |\nabla[(u + \delta)^{\frac{\alpha + (p-1)}{p}}]|^p \phi \, dx \\ & \leq \lambda \int_{\Omega} (u + \delta)^{\alpha+s} \phi \, dx + \frac{1}{p |\alpha|^{p-1}} \int_{\Omega} (u + \delta)^{\alpha+(p-1)} \frac{|\nabla \phi|^p}{\phi^{p-1}} \, dx. \end{aligned}$$

If $\alpha = -(p - 1)$, for any $\delta > 0$, we have the Caccioppoli estimate

$$(3) \quad \frac{(p - 1)^2}{p} \int_{\Omega} |\nabla \log(u + \delta)|^p \phi \, dx + \lambda \int_{\Omega} \frac{u^s}{(u + \delta)^{p-1}} \phi \, dx \leq \frac{1}{p} \int_{\Omega} \frac{|\nabla \phi|^p}{\phi^{p-1}} \, dx.$$

In addition for any $\alpha < 0$ and $\delta > 0$

$$(4) \quad \begin{aligned} & \frac{|\alpha|}{p} \left| \frac{p}{\alpha + (p-1)} \right|^p \int_{\Omega} |\nabla[(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi \, dx \\ & \leq \frac{1}{p |\alpha|^{p-1}} \int_{\Omega} (u + \delta)^{\alpha+(p-1)} \frac{|\nabla \phi|^p}{\phi^{p-1}} \, dx. \end{aligned}$$

Remark 2.1. We underline that when $\alpha > -(p-1)$ we can let $\delta \rightarrow 0^+$ in the energy estimates (2) and (4) to get

$$(5) \quad \begin{aligned} & \frac{p^{p-1} |\alpha|}{[\alpha + (p-1)]^p} \int_{\Omega} |\nabla(u^{\frac{\alpha+(p-1)}{p}})|^p \phi \, dx \\ & \leq \lambda \int_{\Omega} u^{\alpha+s} \phi \, dx + \frac{1}{p |\alpha|^{p-1}} \int_{\Omega} u^{\alpha+(p-1)} \frac{|\nabla \phi|^p}{\phi^{p-1}} \, dx \end{aligned}$$

and for $-(p-1) < \alpha < 0$,

$$(6) \quad \frac{p^{p-1} |\alpha|}{[\alpha + (p-1)]^p} \int_{\Omega} |\nabla(u^{\frac{\alpha+(p-1)}{p}})|^p \phi \, dx \leq \frac{1}{p |\alpha|^{p-1}} \int_{\Omega} u^{\alpha+(p-1)} \frac{|\nabla \phi|^p}{\phi^{p-1}} \, dx.$$

Proof of Lemma 2.1. Let $0 < \phi \in C_c^1(\Omega)$ and $\delta > 0$. Multiply the equation by $(u + \delta)^\alpha \phi$, $\alpha \neq -(p-1)$ and integrate by parts on Ω to get

$$(7) \quad \begin{aligned} & \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \phi (u + \delta)^\alpha \, dx + \alpha \int_{\Omega} |\nabla u|^p (u + \delta)^{\alpha-1} \phi \, dx \\ & = \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla [(u + \delta)^\alpha \phi] \, dx = - \int_{\Omega} A_p u (u + \delta)^\alpha \phi \, dx = \lambda \int_{\Omega} u^\alpha (u + \delta)^\alpha \phi \, dx. \end{aligned}$$

So, for any $\alpha \neq -(p-1)$, we have

$$(8) \quad \begin{aligned} & \frac{|\alpha| p^p}{[\alpha + (p-1)]^p} \int_{\Omega} |\nabla[(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi \, dx = |\alpha| \int_{\Omega} |\nabla u|^p (u + \delta)^{\alpha-1} \phi \, dx \\ & \leq \lambda \int_{\Omega} u^\alpha (u + \delta)^\alpha \phi \, dx \\ & \quad + \int_{\Omega} |\nabla u|^{p-1} (u + \delta)^\alpha |\nabla \phi| \, dx. \end{aligned}$$

Now applying the inequality (79) with $\sigma = \frac{p}{p-1} > 1$ to the second term in the right

hand side of (8), we obtain

$$\begin{aligned} \int_{\Omega} |\nabla u|^{p-1} |\nabla \phi| (u + \delta)^{\alpha} dx &\leq \frac{\varepsilon(p-1)}{p} \left| \frac{p}{\alpha + (p-1)} \right|^p \\ &\quad \times \int_{\Omega} |\nabla [(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi dx \\ &\quad + \frac{1}{p \varepsilon^{p-1}} \int_{\Omega} (u + \delta)^{\alpha+(p-1)} \frac{|\nabla \phi|^p}{\phi^{p-1}} dx. \end{aligned}$$

Simplifying and choosing $\varepsilon = |\alpha|$ such that

$$|\alpha| - \frac{\varepsilon(p-1)}{p} > 0,$$

we arrive at the following energy estimate

$$\begin{aligned} (9) \quad &\frac{|\alpha|}{p} \left| \frac{p}{\alpha + (p-1)} \right|^p \int_{\Omega} |\nabla [(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi dx \\ &\leq \lambda \int_{\Omega} (u + \delta)^{\alpha+s} \phi dx + \frac{1}{p |\alpha|^{p-1}} \int_{\Omega} (u + \delta)^{\alpha+(p-1)} \frac{|\nabla \phi|^p}{\phi^{p-1}} dx. \end{aligned}$$

In the particular case $\alpha < 0$, since u is assumed to be nonnegative, we get from (7),

$$|\alpha| \int_{\Omega} |\nabla u|^p (u + \delta)^{\alpha-1} \phi dx \leq \int_{\Omega} |\nabla u|^{p-1} |\nabla \phi| (u + \delta)^{\alpha} dx.$$

So, proceeding as above, we arrive at

$$\begin{aligned} &\frac{|\alpha|}{p} \left(\frac{p}{|\alpha + (p-1)|} \right)^p \int_{\Omega} |\nabla [(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi dx \\ &\leq \frac{1}{p |\alpha|^{p-1}} \int_{\Omega} (u + \delta)^{\alpha+(p-1)} \frac{|\nabla \phi|^p}{\phi^{p-1}} dx. \end{aligned}$$

Now let us consider the case $\alpha = -(p-1)$, as before, multiplying the equation by $(u + \delta)^{-(p-1)} \phi$, $\delta > 0$, and integrating by parts on Ω , we get

$$\begin{aligned} &\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \frac{\nabla \phi}{(u + \delta)^{p-1}} dx - (p-1) \int_{\Omega} \frac{|\nabla u|^p}{(u + \delta)^p} \phi dx \\ &= \int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla [(u + \delta)^{-(p-1)} \phi] dx \\ &= - \int_{\Omega} \Delta_p u (u + \delta)^{-(p-1)} \phi dx = \lambda \int_{\Omega} u^s (u + \delta)^{-(p-1)} \phi dx. \end{aligned}$$

So

$$(p - 1) \int_{\Omega} \frac{|\nabla u|^p}{(u + \delta)^p} \phi \, dx + \lambda \int_{\Omega} \frac{u^s}{(u + \delta)^{p-1}} \phi \, dx \leq \int_{\Omega} |\nabla u|^{p-1} \frac{|\nabla \phi|}{(u + \delta)^{p-1}} \, dx.$$

Applying Young inequality to the last term of the previous inequality and rewriting

$$\int_{\Omega} \frac{|\nabla u|^p}{(u + \delta)^p} \phi \, dx = \int_{\Omega} |\nabla \log(u + \delta)|^p \phi \, dx$$

we get (3). □

Now we can compute some useful calculations in order to get an explicit expression for all the constants.

Lemma 2.2 (A test function). *Fix two balls $B_{\rho_1} \subset B_{\rho_0} \subset\subset \Omega$. Then there exists a test function $\phi \in C_c^1(\Omega)$ which is radially symmetric and piecewise C^2 such that $\text{supp}(\phi) = B_{\rho_0}$, $\phi = 1$ on B_{ρ_1} , and satisfies*

$$(10) \quad \frac{|\nabla \phi|^p}{\phi^{p-1}} \leq \frac{2^{p-1} p^p}{(\rho_0 - \rho_1)^p} \quad \text{and} \quad \|\nabla \phi\|_{\infty} \leq \frac{p}{\rho_0 - \rho_1} \quad \text{for any } p > 1.$$

Proof. Consider the radial test function defined on $B_{\rho_0} \subset\subset \Omega$

$$\phi(|x|) = \begin{cases} 1, & \text{if } 0 \leq |x| \leq \rho_1 \\ 1 - \frac{2^{p-1}(|x| - \rho_1)^p}{(\rho_0 - \rho_1)^p}, & \text{if } \rho_1 < |x| \leq \frac{\rho_0 + \rho_1}{2} \\ \frac{2^{p-1}(\rho_0 - |x|)^p}{(\rho_0 - \rho_1)^p}, & \text{if } \frac{\rho_0 + \rho_1}{2} < |x| \leq \rho_0 \\ 0, & \text{if } |x| > \rho_0 \end{cases}$$

for any $0 < \rho_1 < \rho_0$. We have

$$\nabla \phi(|x|) = \begin{cases} 0, & \text{if } 0 \leq |x| \leq \rho_1 \text{ or if } |x| > \rho_0 \\ -\frac{2^{p-1} p (|x| - \rho_1)^{p-1}}{(\rho_0 - \rho_1)^p} \frac{x}{|x|}, & \text{if } \rho_1 < |x| \leq \frac{\rho_0 + \rho_1}{2} \\ -\frac{2^{p-1} p (\rho_0 - |x|)^{p-1}}{(\rho_0 - \rho_1)^p} \frac{x}{|x|}, & \text{if } \frac{\rho_0 + \rho_1}{2} < |x| \leq \rho_0. \end{cases}$$

So we easily obtain the bounds (10). □

Remark 2.2. As a consequence of the first inequality in (10), we have that

$$\int_{\Omega} \frac{|\nabla \phi|^p}{\phi^{p-1}} \, dx \leq \frac{2^{p-1} p^p |B_{\rho_0}|}{(\rho_0 - \rho_1)^p} = \frac{2^{p-1} p^p \rho_0^d \omega_d}{(\rho_0 - \rho_1)^p}.$$

Corollary 2.1 (Quantitative Caccioppoli Estimates). *Let $\delta > 0$. Under the assumption of Lemma 2.1 and using the test function ϕ of Lemma 2.2, we have the quantitative Caccioppoli estimates, for any $\delta > 0$:*

$$(11) \quad \frac{(p-1)^2}{p} \int_{B_{\rho_1}} |\nabla \log(u + \delta)|^p \, dx + \lambda \int_{B_{\rho_1}} \frac{u^s}{(u + \delta)^{p-1}} \, dx \leq \frac{p^{p-1} 2^{p-1} \rho_0^d \omega_d}{(\rho_0 - \rho_1)^p}.$$

Proof. Using (3) with ϕ as in Lemma 2.2 and recalling Remark 2.2 we easily obtain the desired result. □

Remark 2.3. Letting $\delta \rightarrow 0^+$ in (11), we get

$$(12) \quad \lambda \int_{B_{\rho_1}} u^{s-(p-1)} \, dx \leq \frac{p^{p-1} 2^{p-1} \omega_d \rho_0^d}{(\rho_0 - \rho_1)^p}.$$

As a consequence of this fact in Section 7 we obtain a local absolute upper bound in the range $p - 1 < s < s_c^* := r(p - 1)/p$, r defined in (14) below and a local absolute lower bound if $u \not\equiv 0$ on B_{ρ_0} and $0 \leq s < p - 1$.

3 - Sobolev and reverse Poincaré inequalities

In this section we will recall the Sobolev inequalities that will be used throughout the paper and we also show how they combine with the energy inequalities of the previous section to give a kind of reverse Poincaré inequalities, that will be necessary for the upper bounds when dealing with unbounded coefficients.

Sobolev inequalities. Our local bounds will be a consequence of the Sobolev imbedding theorems on balls $B_\rho \subset \mathbb{R}^d$. Indeed the following Sobolev type inequalities hold true:

$$(13) \quad \|g\|_{L^r(B_\rho)}^p \leq \mathcal{S}_p^p \left(\|\nabla g\|_{L^p(B_\rho)}^p + \frac{1}{\rho^p} \|g\|_{L^p(B_\rho)}^p \right)$$

for any $g \in W^{1,p}(\Omega)$, where Ω is a bounded open domain of \mathbb{R}^d with smooth boundary,

$B_\rho \subset \Omega$, and

$$(14) \quad \begin{cases} \text{if } p < d, & r = p^* = \frac{pd}{d-p}, & \mathcal{S}_p \\ \text{if } p = d, & r \in (p, +\infty), & \mathcal{S}_p = \mathcal{S}'_p \text{diam}(\Omega)^{\frac{d}{r}} \\ \text{if } p > d, & r = +\infty, & \mathcal{S}_p = \mathcal{S}'_p \text{diam}(\Omega)^{1-\frac{d}{p}} \end{cases}$$

and $\mathcal{S}_p, \mathcal{S}'_p > 0$ only depends on p, d , see e.g. Theorem 3.11 and 3.12 of [34]. On the other hand, whenever $g \in W_0^{1,p}(B_\rho)$ we have

$$(15) \quad \|g\|_{L^r(B_\rho)}^p \leq \mathcal{S}_p^p \|\nabla g\|_{L^p(B_\rho)}^p,$$

where r is defined in (14) and \mathcal{S}_p is the Sobolev constant, which only depends on p, d , see e.g. Theorem 3.9 of [34]. We will denote by r the Sobolev exponent corresponding to $W_0^{1,p}(B_\rho)$ through all the paper.

Now we state and prove a lemma originally due to Trudinger [53] (see Lemma 5.1 p. 745 there). For a proof in the case $p = 2$ see Lemma 3.2 in [4].

Lemma 3.1. *Let $v \in L^r(B_\rho)$ and $b \in L^m(B_\rho)$ for $m > r/(r - p)$ with r defined in (14). Then for any $\gamma > 0$ the following inequality holds*

$$(16) \quad \int_{B_\rho} b v^p \, dx \leq \gamma \left(\int_{B_\rho} v^r \, dx \right)^{\frac{p}{r}} + \frac{K_{m,p,r}}{\gamma^{\frac{mp+r}{m(r-p)-r}}} |B_\rho|^{\frac{p}{r}} \left(\int_{B_\rho} b^m \, dx \right)^{\frac{r}{m(r-p)-r}} \int_{B_\rho} v^p \, dx,$$

where

$$(17) \quad K_{m,p,r} := \frac{m(r-p) - r}{rm} \left(\frac{pm+r}{rm} \right)^{\frac{mp+r}{m(r-p)-r}}.$$

Proof. Let us estimate for any $0 < \gamma_1 < p$,

$$\begin{aligned} \int_{B_\rho} b v^{(p-\gamma_1)+\gamma_1} \, dx &\leq_{(a)} \left(\int_{B_\rho} v^{(p-\gamma_1)\frac{r}{p}} \, dx \right)^{\frac{p}{r}} \left(\int_{B_\rho} b^{\frac{r}{r-p}} v^{\gamma_1 \frac{r}{r-p}} \, dx \right)^{\frac{r-p}{r}} \\ &\leq_{(b)} \left(\int_{B_\rho} v^r \, dx \right)^{\frac{p-\gamma_1}{r}} |B_\rho|^{\frac{\gamma_1}{r}} \left(\int_{B_\rho} b^{\frac{r}{r-p}} v^{\gamma_1 \frac{r}{r-p}} \, dx \right)^{\frac{r-p}{r}} \end{aligned}$$

$$\begin{aligned}
&\leq_{(c)} \frac{\gamma_0(p - \gamma_1)}{p} \left(\int_{B_\rho} v^r \, dx \right)^{\frac{p}{r}} \\
&\quad + \frac{\gamma_1}{p} \frac{1}{\gamma_0^{\frac{p-\gamma_1}{\gamma_1}}} |B_\rho|^{\frac{p}{r}} \left(\int_{B_\rho} b^{\frac{r}{r-p}} v^{\gamma_1 \frac{r}{r-p}} \, dx \right)^{\frac{p(r-p)}{\gamma_1 r}} \\
&\leq_{(d)} \frac{\gamma_0(p - \gamma_1)}{p} \left(\int_{B_\rho} v^r \, dx \right)^{\frac{p}{r}} + \frac{\gamma_1}{p} \frac{1}{\gamma_0^{\frac{p-\gamma_1}{\gamma_1}}} |B_\rho|^{\frac{p}{r}} \\
&\quad \times \left(\int_{B_\rho} b^m \, dx \right)^{\frac{p}{\gamma_1 m}} \left(\int_{B_\rho} v^{\frac{\gamma_1 r m}{m(r-p)-r}} \, dx \right)^{\frac{p[m(r-p)-r]}{\gamma_1 r m}} \\
&=_{(e)} \frac{(pm + r)\gamma_0}{rm} \left(\int_{B_\rho} v^r \, dx \right)^{\frac{p}{r}} + \frac{m(r-p) - r}{mr} \\
&\quad \times \frac{1}{\gamma_0^{\frac{mp+r}{m(r-p)-r}}} |B_\rho|^{\frac{p}{r}} \left(\int_{B_\rho} b^m \, dx \right)^{\frac{r}{m(r-p)-r}} \int_{B_\rho} v^p \, dx,
\end{aligned}$$

where in (a) we have used Hölder inequality with exponents r/p and $r/(r-p)$, in (b) with $p/(p-\gamma_1)$ and p/γ_1 ; in (c) we have applied the Young inequality (79), with $\varepsilon = \gamma_0$, $\sigma = p/(p-\gamma_1)$. In (d) we have used again Hölder inequality with exponents $m(r-p)/r > 1$, since we are assuming $m > r/(r-p)$, and $m(r-p)/[m(r-p)-r]$ and in (e) we have put

$$0 < \gamma_1 = \frac{p[m(r-p) - r]}{rm} < p.$$

To obtain the desired result it is sufficient to take

$$\gamma = \gamma_0 \frac{pm + r}{rm}.$$

□

Theorem 3.1 (Reverse Poincaré inequality). *Let u be a weak solution to $-\Delta_p u = b u^{p-1}$ in B_ρ , with $p > 1$. Let $b \in L^m(B_\rho)$, for $m > r/(r-p)$ and r be defined in (14). Suppose that $u \in L^{x+(p-1)}(B_\rho)$. Then for any positive test function*

$1 \geq \phi \in C_c^1(B_\rho)$ and any $\alpha > 0$ the following estimate holds true

$$\int_{B_\rho} \left| \nabla \left(u^{\frac{\alpha+(p-1)}{p}} \right) \right|^p \phi^p \, dx \leq A(b) \int_{B_\rho} u^{\alpha+(p-1)} \, dx,$$

with

$$\begin{aligned} A(b) := & \left\{ \left[1 + 2 \left(\frac{\alpha + (p-1)}{\alpha} \right)^p \right] \|\nabla \phi\|_\infty^p \right. \\ & + K_{m,p,r} \frac{2p}{\alpha} \left(\frac{2^p p S_p^p}{\alpha} \right)^{\frac{mp+r}{m(r-p)-r}} \left(\frac{\alpha + (p-1)}{p} \right)^{\frac{mpr}{m(r-p)-r}} \\ & \left. \times |B_\rho|^{\frac{p}{r}} \left(\int_{\tilde{B}_\rho} b^m \, dx \right)^{\frac{r}{m(r-p)-r}} \right\}, \end{aligned}$$

$K_{m,p,r}$ given in (17).

Remark 3.1. We underline that the requirement $u \in L^{\alpha+(p-1)}(B_\rho)$ will be dispensed later, without further comment by using a Moser iteration technique.

Proof. We divide the proof in few steps.

• **STEP 1. Energy estimates.** Multiplying $-\Delta_p u = b u^{p-1}$ by $(u + \delta)^\alpha \phi^p$, $1 \geq \phi \in C_c^1(B_\rho)$, $\alpha > 0$, $\delta > 0$ and integrating by parts on B_ρ , we get

$$\begin{aligned} (18) \quad \alpha \int_{\tilde{B}_\rho} |\nabla u|^p (u + \delta)^{\alpha-1} \phi^p \, dx & \leq p \int_{\tilde{B}_\rho} |\nabla u|^{p-1} |\nabla \phi| \phi^{p-1} (u + \delta)^\alpha \, dx \\ & + \int_{\tilde{B}_\rho} b u^{p-1} (u + \delta)^\alpha \phi^p \, dx. \end{aligned}$$

Using (79) with $\sigma = p/(p-1)$ to estimate the first term in the right hand side of (18), we obtain

$$\begin{aligned} & [\alpha - \varepsilon(p-1)] \left(\frac{p}{\alpha + (p-1)} \right)^p \int_{\tilde{B}_\rho} |\nabla [(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi^p \, dx \\ & \leq \int_{\tilde{B}_\rho} b (u + \delta)^{\alpha+(p-1)} \phi^p \, dx + \frac{1}{\varepsilon^{p-1}} \int_{\tilde{B}_\rho} (u + \delta)^{\alpha+(p-1)} |\nabla \phi|^p \, dx. \end{aligned}$$

Choosing $\varepsilon = \alpha/p$, we arrive at

$$(19) \quad \begin{aligned} & \frac{\alpha}{p} \left(\frac{p}{\alpha + (p-1)} \right)^p \int_{B_\rho} |\nabla[(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi^p \, dx \\ & \leq \int_{B_\rho} b(u + \delta)^{\alpha+(p-1)} \phi^p \, dx + \frac{p^{p-1}}{\alpha^{p-1}} \int_{B_\rho} (u + \delta)^{\alpha+(p-1)} |\nabla \phi|^p \, dx. \end{aligned}$$

• **STEP 2. Sobolev inequality in $W_0^{1,p}(B_\rho)$.** We apply inequality (16) to $v = (u + \delta)^{\frac{\alpha+(p-1)}{p}} \phi \in W_0^{1,p}(B_\rho)$, so that for any $\gamma > 0$:

$$(20) \quad \begin{aligned} \int_{B_\rho} b(u + \delta)^{\alpha+(p-1)} \phi^p \, dx & \leq \gamma \left(\int_{B_\rho} (u + \delta)^{[\alpha+(p-1)]\frac{p}{r}} \phi^r \, dx \right)^{\frac{p}{r}} \\ & + \frac{K_{m,p,r}}{\gamma^{\frac{m(p+r)}{m(r-p)-r}}} |B_\rho|^{\frac{p}{r}} \left(\int_{B_\rho} b^m \, dx \right)^{\frac{r}{m(r-p)-r}} \\ & \times \int_{B_\rho} (u + \delta)^{\alpha+(p-1)} \phi^p \, dx, \end{aligned}$$

where $K_{m,p,r}$ is given in (17). Since $v = (u + \delta)^{\frac{\alpha+(p-1)}{p}} \phi \in W_0^{1,p}(B_\rho)$, the Sobolev inequality (15) reads

$$\begin{aligned} \left(\int_{B_\rho} [(u + \delta)^{\frac{\alpha+(p-1)}{p}} \phi]^r \, dx \right)^{\frac{p}{r}} & \leq S_p^p 2^{p-1} \int_{B_\rho} |\nabla[(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi^p \, dx \\ & + S_p^p 2^{p-1} \int_{B_\rho} (u + \delta)^{\alpha+(p-1)} |\nabla \phi|^p \, dx. \end{aligned}$$

We combine the above Sobolev inequality with (20) to get

$$(21) \quad \begin{aligned} \int_{B_\rho} b(u + \delta)^{\alpha+(p-1)} \phi^p \, dx & \leq \gamma S_p^p 2^{p-1} \int_{B_\rho} |\nabla[(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi^p \, dx \\ & + \gamma S_p^p 2^{p-1} \int_{B_\rho} (u + \delta)^{\alpha+(p-1)} |\nabla \phi|^p \, dx \\ & + \frac{K_{m,p,r}}{\gamma^{\frac{m(p+r)}{m(r-p)-r}}} |B_\rho|^{\frac{p}{r}} \left(\int_{B_\rho} b^m \, dx \right)^{\frac{r}{m(r-p)-r}} \\ & \times \int_{B_\rho} (u + \delta)^{\alpha+(p-1)} \phi^p \, dx. \end{aligned}$$

- STEP 3. Putting together (19) and (21) and recalling that $\phi \leq 1$, we obtain

$$\begin{aligned} & \left[\frac{\alpha}{p} \left(\frac{p}{\alpha + (p-1)} \right)^p - \gamma S_p^p 2^{p-1} \right] \int_{B_\rho} |\nabla[(u + \delta)^{\frac{\alpha+(p-1)}{p}}]|^p \phi^p \, dx \\ & \leq \left\{ \left[\gamma S_p^p 2^{p-1} + \frac{p^{p-1}}{\alpha^{p-1}} \|\nabla \phi\|_\infty^p + \frac{K_{m,p,r}}{\gamma^{\frac{mp+r}{m(r-p)-r}}} |B_\rho|^{\frac{p}{r}} \left(\int_{B_\rho} b^m \, dx \right)^{\frac{r}{m(r-p)-r}} \right] \right. \\ & \quad \left. \times \int_{B_\rho} (u + \delta)^{\alpha+(p-1)} \, dx. \right. \end{aligned}$$

Choosing

$$\gamma = \left(\frac{p}{\alpha + (p-1)} \right)^p \frac{\alpha}{p 2^p S_p^p}$$

and letting $\delta \rightarrow 0$ we obtain the desired result. □

Remark 3.2. If we take ϕ as in Lemma 2.2 we obtain

$$\int_{B_{\rho_1}} |\nabla(u^{\frac{\alpha+(p-1)}{p}})|^p \, dx \leq A(b) \int_{B_{\rho_0}} u^{\alpha+(p-1)} \, dx,$$

with

$$\begin{aligned} (22) \quad A(b) := & \left\{ \left[1 + 2 \left(\frac{\alpha + (p-1)}{\alpha} \right)^p \right] \frac{p^p}{(\rho_0 - \rho_1)^p} + K_{m,r,p} \left(\frac{2^p p S_p^p}{\alpha} \right)^{\frac{mp+r}{m(r-p)-r}} \right. \\ & \left. \times \frac{2p}{\alpha} \left(\frac{\alpha + (p-1)}{p} \right)^{\frac{mpr}{m(r-p)-r}} \omega_d^{\frac{p}{r}} \rho_0^{\frac{dp}{r}} \left(\int_{B_{\rho_0}} b^m \, dx \right)^{\frac{r}{m(r-p)-r}} \right\}, \end{aligned}$$

for all $\alpha > 0$, $m > r/(r-p)$, and $K_{m,p,r}$ as in (17).

4 - Local upper bounds

This section is devoted to the proof of quantitative local upper bounds for local nonnegative weak solutions to $-\Delta_p u = \lambda u^s$, for any $\lambda \geq 0$ and any $s \geq 0$. We also get quantitative local estimates for solutions to the problem $-\Delta_p u = b(x)u^{p-1}$ with

$b \in L^m$, eventually unbounded. We prove our results for nonnegative solutions, but the careful reader can realize that almost the same proof holds for nonnegative subsolutions, or for solutions with any sign.

4.1 - Local upper bounds I. The upper Moser iteration

The first form of the upper bounds that we present in this section, is a consequence of energy estimates, Caccioppoli inequalities and the “local” Sobolev inequality on balls.

Theorem 4.1 (Local Upper Estimates). *Let $\Omega \subset \mathbb{R}^d$ and $\lambda \geq 0$. Let u be a local nonnegative weak solution to $-\Delta_p u = \lambda u^s$ in Ω with $p > 1$, $0 \leq s < r - 1$ and r as in (14). Then the following bound holds true for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$ and for any $q > \bar{q} := [s - (p - 1)]_+ r / (r - p)$*

$$(23) \quad \|u\|_{\infty, R_\infty} \leq I_{\infty, q} \left(\int_{B_{R_0}} u^q \, dx \right)^{\frac{(r-p)\mu}{r}} \left(\int_{B_{R_\infty}} u^{[s-(p-1)]_+} \, dx \right)^{-\mu}$$

where $\mu = r / \{(r - p)q - r[s - (p - 1)]_+\}$ and the constant $I_{\infty, q}$ depends on $d, p, s, q, r, R_0, R_\infty$, and when $s \neq p - 1$ does not depend on λ , see an explicit expression in formula (26) below.

Moreover, when $0 \leq s \leq p - 1$, the above estimate takes the simplified form:

$$(24) \quad \|u\|_{\infty, R_\infty} \leq I_{\infty, q} \left(\int_{B_{R_0}} u^q \, dx \right)^{\frac{1}{q}},$$

and holds for all $q > 0$. The constant $I_{\infty, q}$ is the same as above and is given in formula (26) below.

Remark on the result. Inequality (23) is a kind of reverse Hölder inequality, indeed we can rewrite it as:

$$(25) \quad \|u\|_{[s-(p-1)]_+, R_\infty}^{\mu[s-(p-1)]_+} \|u\|_{\infty, R_\infty} \leq C \|u\|_{q, R_0}^{\frac{q(r-p)\mu}{r}}.$$

This form makes clearer the fact that if there is a constant that makes true (25) for a $q \geq \bar{q}$, then by Hölder inequality, the same inequality holds true for all $q' > q$, with the same constant. The same applies to (23).

Remark on the constant. The proof below allows us to find an explicit expression of the constant:

$$(26) \quad I_{\infty,q} = \left[\frac{|B_{R_0}|^{\frac{r-p}{r}+1}}{|B_{R_\infty}|} \frac{S_p^p q^p p^p c_1}{r^p (R_0 - R_\infty)^p} \left\{ \frac{A_{s,0} r}{p q} \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \right. \right. \\ \left. \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p c_2 + \frac{c_1^{p-1} 2^{p-1}}{c_0^p} \right\} \left(\frac{r}{p} \right)^{\frac{pr}{r-p}} \right]^{\frac{r}{q(r-p) - r[s-(p-1)]_+}}$$

with

$$\frac{1}{c_0^p} \leq \left(\frac{r}{(r-p)q - [s-(p-1)]_+ r} \right)^p,$$

$$c_1 := \begin{cases} \frac{pq}{pq - r(p-1)}, & \text{if } q > s_c^* = \frac{r(p-1)}{p}, \\ \max_{i=0,1} \left[\frac{\left(\frac{r}{p}\right)^{j_0+i-1} q}{\left(\frac{r}{p}\right)^{j_0+i-1} q - (p-1)} \right], & \text{if } 0 < q < s_c^* = \frac{r(p-1)}{p}, \end{cases}$$

$$c_2 := \max \left\{ \frac{|pq - r(p-1)| r^p}{(pq)^{p+1}}, \left(\frac{p}{p-1} \right)^p \frac{1}{(p+1)^{p+1}} \right\}$$

and

$$(27) \quad A_{s,0} := \begin{cases} \frac{\lambda}{p^{p-1}} R_0^p, & \text{if } s = p-1, \\ 2^{p-1+d}, & \text{if } s \neq p-1. \end{cases}$$

Moreover when $0 < q < s_c^*$ we require the additional condition

$$(28) \quad j_0 := i.p. \left[\frac{\log \left(\frac{r(p-1)(r-p) - rp[s-(p-1)]_+}{qp(r-p) - rp[s-(p-1)]_+} \right)}{\log \frac{r}{p}} \right] \neq \frac{\log \left(\frac{r(p-1)(r-p) - rp[s-(p-1)]_+}{qp(r-p) - rp[s-(p-1)]_+} \right)}{\log \frac{r}{p}}$$

where $i.p.[t]$ denotes the integer part of $t \in \mathbb{R}$.

Finally, we would like to remark that this latter condition (28) is not really essential: indeed, we can obtain an explicit constant $I_{\infty,q}$ for a $q > \bar{q}$ such that

$$i.p. \left[\frac{\log \left(\frac{r(p-1)(r-p) - rp[s-(p-1)]_+}{qp(r-p) - rp[s-(p-1)]_+} \right)}{\log \frac{r}{p}} \right] = \frac{\log \left(\frac{r(p-1)(r-p) - rp[s-(p-1)]_+}{qp(r-p) - rp[s-(p-1)]_+} \right)}{\log \frac{r}{p}},$$

simply by considering a $q' \in (\bar{q}, q)$ such that condition (28) holds so that the explicit constant is given by $I_{\infty,q'}$; then by the remark after formula (25), we obtain the desired bound also for $q > q'$ with the same constant ($I_{\infty,q} = I_{\infty,q'}$) as a consequence of Hölder inequality.

Proof of Theorem 4.1. We are going to use the energy estimate (2) for any $\alpha > -(p - 1)$, $\alpha \neq 0$, to prove $L^q - L^\infty$ local estimates via Moser iteration, keeping track all the constants. We underline, as in Remark 3.1, that the requirement $u \in L^{\alpha+(p-1)}(B_\rho)$ needed at each iteration step, will be dispensed by the previous one. In particular, notice that $u \in L^{\alpha+(p-1)}(B_\rho)$ implies $g = u^{[\alpha+(p-1)]/p} \in W_{loc}^{1,p}(\Omega)$ by the energy estimates (5). Therefore, we only have to ensure the integrability condition at the first step, which gives rise to condition $q > \bar{q} := [s - (p - 1)]_+ r / (r - p)$.

We divide the proof in several steps.

- **STEP 1.** Let ϕ be the function defined in Lemma 2.2. The local Sobolev inequality (13) on the ball B_{ρ_1} , where ρ_1 is any real positive number such that $B_{\rho_1} \subset B_{\rho_0} \subset\subset \Omega$, applied to $g = u^{[\alpha+(p-1)]/p}$ together with the energy estimate (5), gives, using the properties of the function ϕ , established in Lemma 2.2,

$$\begin{aligned}
 \left[\int_{B_{\rho_1}} u^{[\alpha+(p-1)]\frac{p}{p}} dx \right]^{\frac{p}{r}} &\leq S_p^p \left[\int_{B_{\rho_1}} |\nabla(u^{\frac{\alpha+(p-1)}{p}})|^p dx \right. \\
 &\quad \left. + \frac{1}{\rho_1^p} \int_{B_{\rho_0}} u^{\alpha+(p-1)} dx \right] \\
 (29) \quad &\leq S_p^p \left\{ \frac{p}{|\alpha|} \left(\frac{\alpha + (p - 1)}{p} \right)^p \lambda \int_{B_{\rho_0}} u^{\alpha+s} dx \right. \\
 &\quad \left. + \left[\frac{1}{\rho_1^p} + \left(\frac{\alpha + (p - 1)}{|\alpha|} \right)^p \frac{2^{p-1}}{(\rho_0 - \rho_1)^p} \right] \right. \\
 &\quad \left. \times \int_{B_{\rho_0}} u^{\alpha+(p-1)} dx \right\}.
 \end{aligned}$$

- **STEP 2.** *Caccioppoli estimates and first iteration step.* Now we need to split into two cases, namely $0 \leq s \leq p - 1$ and $p - 1 < s < r - 1$.

Superlinear case: $p - 1 < s < r - 1$. We continue estimate (29) as follows:

$$\begin{aligned}
 \left[\int_{B_{\rho_1}} u^{[\alpha+(p-1)]\frac{p}{r}} dx \right]^{\frac{r}{p}} &\leq S_p^p \left\{ \frac{p}{|\alpha|} \left(\frac{\alpha + (p-1)}{p} \right)^p \lambda \right. \\
 &\quad \left. + \left[\frac{1}{\rho_1^p} + \left(\frac{\alpha + (p-1)}{|\alpha|} \right)^p \frac{2^{p-1}}{(\rho_0 - \rho_1)^p} \right] \right. \\
 &\quad \left. \times \frac{\int_{B_{\rho_0}} u^{\alpha+(p-1)} dx}{\int_{B_{\rho_0}} u^{\alpha+s} dx} \right\} \int_{B_{\rho_0}} u^{\alpha+s} dx \\
 (30) \quad &\leq_{(a)} S_p^p \frac{|B_{\rho_0}|}{\|u\|_{s-(p-1),\rho_0}^{s-(p-1)}} \left\{ \frac{p}{|\alpha|} \left(\frac{\alpha + (p-1)}{p} \right)^p \lambda \right. \\
 &\quad \times \frac{\|u\|_{s-(p-1),\rho_0}^{s-(p-1)}}{|B_{\rho_0}|} \\
 &\quad \left. + \left[\frac{1}{\rho_1^p} + \left(\frac{\alpha + (p-1)}{|\alpha|} \right)^p \frac{2^{p-1}}{(\rho_0 - \rho_1)^p} \right] \right\} \\
 &\quad \times \int_{B_{\rho_0}} u^{\alpha+s} dx \\
 &\leq_{(b)} \frac{S_p^p |B_{\rho_0}|}{(\rho_0 - \rho_1)^p \|u\|_{s-(p-1),\rho_0}^{s-(p-1)}} \\
 &\quad \times \left\{ \frac{[\alpha + (p-1)]^p 2^{p-1+d}}{|\alpha|} \left(\frac{\rho_0 - \rho_1}{\rho_0} \right)^p \right. \\
 &\quad \left. + \left(\frac{\rho_0 - \rho_1}{\rho_1} \right)^p + \frac{[\alpha + (p-1)]^p 2^{p-1}}{|\alpha|^p} \right\} \\
 &\quad \times \int_{B_{\rho_0}} u^{\alpha+s} dx.
 \end{aligned}$$

In (a) we have used the convexity in the variable $t > 0$ of the function $N(t) = \log \|u\|_t^t$. Hence, since the incremental quotient is increasing (see for example [50] for more details), choosing $\alpha + (p - 1) \geq \bar{\alpha} > 0$, we obtain

$$\frac{N(\bar{\alpha} + s - (p - 1)) - N(\bar{\alpha})}{s - (p - 1)} \leq \frac{N(\alpha + s) - N(\alpha + (p - 1))}{s - (p - 1)}$$

that is

$$\frac{\|u\|_{\bar{x}+s-(p-1),\rho_0}^{\bar{x}+s-(p-1)}}{\|u\|_{\bar{x},\rho_0}^{\bar{x}}} \leq \frac{\|u\|_{\alpha+s,\rho_0}^{\alpha+s}}{\|u\|_{\alpha+(p-1),\rho_0}^{\alpha+(p-1)}}.$$

Then, using Hölder inequality with exponents $[\bar{x} + s - (p - 1)]/[s - (p - 1)] > 1$ and $[\bar{x} + s - (p - 1)]/\bar{x} > 1$, since we are assuming $s > p - 1$, we have

$$\begin{aligned} \frac{\|u\|_{\alpha+s,\rho_0}^{\alpha+s}}{\|u\|_{\alpha+(p-1),\rho_0}^{\alpha+(p-1)}} &\geq \frac{\|u\|_{\bar{x}+s-(p-1),\rho_0}^{\bar{x}+s-(p-1)}}{\|u\|_{\bar{x},\rho_0}^{\bar{x}}} = \frac{\|u\|_{\bar{x}+s-(p-1),\rho_0}^{\bar{x}}}{\|u\|_{\bar{x},\rho_0}^{\bar{x}}} \|u\|_{\bar{x}+s-(p-1),\rho_0}^{s-(p-1)} \\ &\geq |B_{\rho_0}|^{-\frac{s-(p-1)}{\bar{x}+s-(p-1)}} \|u\|_{s-(p-1),\rho_0}^{s-(p-1)} |B_{\rho_0}|^{-\frac{\bar{x}}{\bar{x}+s-(p-1)}} \\ &= \frac{\|u\|_{s-(p-1),\rho_0}^{s-(p-1)}}{|B_{\rho_0}|}. \end{aligned}$$

In (b) we have used the Caccioppoli estimate (12) with ρ_0 and $2\rho_0$, that is

$$(31) \quad \frac{\lambda \|u\|_{s-(p-1),\rho_0}^{s-(p-1)}}{|B_{\rho_0}|} \leq \frac{p^{p-1} 2^{p-1}}{(2\rho_0 - \rho_0)^p} \frac{|B_{2\rho_0}|}{|B_{\rho_0}|} = \frac{p^{p-1} 2^{p-1+d}}{\rho_0^p}.$$

Sublinear case: $0 \leq s \leq p - 1$. We first assume $0 \leq s < p - 1$, we discuss the case $s = p - 1$ separately. We continue estimate (29) as follows:

$$\begin{aligned} (32) \quad \left[\int_{B_{\rho_1}} u^{[\alpha+(p-1)]_p^+} dx \right]^{\frac{p}{r}} &\leq S_p^p \left\{ \frac{p}{|\alpha|} \left(\frac{\alpha + (p - 1)}{p} \right)^p \lambda \frac{\int_{B_{\rho_0}} u^{\alpha+s} dx}{\int_{B_{\rho_0}} u^{\alpha+(p-1)} dx} \right. \\ &\quad \left. + \frac{1}{\rho_1^p} + \frac{[\alpha + (p - 1)]^p}{|\alpha|^p} \frac{2^{p-1}}{(\rho_0 - \rho_1)^p} \right\} \\ &\quad \times \int_{B_{\rho_0}} u^{\alpha+(p-1)} dx \\ &\leq \frac{S_p^p}{(\rho_0 - \rho_1)^p} \left\{ \frac{[\alpha + (p - 1)]^p 2^{p-1+d}}{|\alpha|} \left(\frac{\rho_0 - \rho_1}{\rho_0} \right)^p \right. \\ &\quad \left. + \left(\frac{\rho_0 - \rho_1}{\rho_1} \right)^p + \frac{[\alpha + (p - 1)]^p 2^{p-1}}{|\alpha|^p} \right\} \\ &\quad \times \int_{B_{\rho_0}} u^{\alpha+(p-1)} dx. \end{aligned}$$

Indeed the properties of the function $N(t) = \log \|u\|_t^t$ give

$$\frac{N(\bar{x}) - N(\bar{x} + s - (p - 1))}{(p - 1) - s} \leq \frac{N(\alpha + (p - 1)) - N(\alpha + s)}{(p - 1) - s}$$

that is

$$\frac{\|u\|_{\alpha+s, \rho_0}^{\alpha+s}}{\|u\|_{\alpha+(p-1), \rho_0}^{\alpha+(p-1)}} \leq \frac{\|u\|_{\bar{x}+s-(p-1), \rho_0}^{\bar{x}+s-(p-1)}}{\|u\|_{\bar{x}, \rho_0}^{\bar{x}}}.$$

Then, using Hölder inequality, the following reverse Hölder inequality

$$\left(\int_{B_{\rho_0}} u^{\bar{x}} \, dx \right)^{-\frac{(p-1)-s}{\bar{x}}} \leq \frac{\int_{B_{\rho_0}} u^{s-(p-1)} \, dx}{|B_{\rho_0}|^{\frac{\bar{x}+(p-1)-s}{\bar{x}}}}$$

holds and (31) gives

$$\begin{aligned} \frac{\|u\|_{\alpha+s, \rho_0}^{\alpha+s}}{\|u\|_{\alpha+(p-1), \rho_0}^{\alpha+(p-1)}} &\leq \frac{\|u\|_{\bar{x}+s-(p-1), \rho_0}^{\bar{x}+s-(p-1)}}{\|u\|_{\bar{x}, \rho_0}^{\bar{x}}} \leq \frac{|B_{\rho_0}|^{\frac{(p-1)-s}{\bar{x}}}}{\|u\|_{\bar{x}, \rho_0}^{(p-1)-s}} \\ &\leq \frac{\int_{B_{\rho_0}} u^{s-(p-1)} \, dx}{|B_{\rho_0}|} \leq \frac{1}{\lambda} \frac{p^{p-1} 2^{p-1+d}}{\rho_0^p}. \end{aligned}$$

Notice that when $s = p - 1$, we obtain from (29) directly

$$\begin{aligned} \left[\int_{B_{\rho_1}} u^{[\alpha+(p-1)]\frac{p}{r}} \, dx \right]^{\frac{r}{p}} &\leq \frac{S_p^p}{(\rho_0 - \rho_1)^p} \left\{ \frac{p}{|\alpha|} \frac{[\alpha + (p - 1)]^p \lambda}{p^p} (\rho_0 - \rho_1)^p \right. \\ (33) \quad &\quad \left. + \left(\frac{\rho_0 - \rho_1}{\rho_1} \right)^p + \frac{[\alpha + (p - 1)]^p 2^{p-1}}{|\alpha|^p} \right\} \\ &\quad \times \int_{B_{\rho_0}} u^{\alpha+(p-1)} \, dx. \end{aligned}$$

• **STEP 3. The first iteration step.** Now we are ready to write the first iteration step for all $s \geq 0$. Let $\beta = \alpha + (p - 1) \geq \beta_0 > 0$ and recall that we are requiring $\beta \neq (p - 1)$ as well, then inequalities (30), (32) and (33) can be written as

$$\left[\int_{B_{\rho_1}} u^{\beta\frac{p}{r}} \, dx \right]^{\frac{r}{p}} \leq I(\beta, s, p, \rho_1, \rho_0) \int_{B_{\rho_0}} u^{\beta+[s-(p-1)]_+} \, dx$$

with

$$(34) \quad I(\beta, s, p, \rho_1, \rho_0) := \frac{S_p^p}{(\rho_0 - \rho_1)^p} \frac{|B_{\rho_0}|}{\int_{B_{\rho_0}} u^{[s-(p-1)]_+} dx} \left\{ \frac{\beta^p A_s}{|\beta - (p-1)|} \right. \\ \left. \times \left(\frac{\rho_0 - \rho_1}{\rho_0} \right)^p + \left(\frac{\rho_0 - \rho_1}{\rho_1} \right)^p + \frac{\beta^p 2^{p-1}}{|\beta - (p-1)|^p} \right\},$$

and

$$(35) \quad A_s := \begin{cases} \frac{\lambda}{p^{p-1}} \rho_0^p, & \text{if } s = p - 1, \\ 2^{p-1+d}, & \text{if } s \neq p - 1. \end{cases}$$

• **STEP 4. The Moser iteration.** Let us define the sequence of exponents $\beta_n > 0$ so that

$$\beta_n + [s - (p - 1)]_+ = \beta_{n-1} \frac{r}{p} \quad \Rightarrow \quad \beta_n = \beta_{n-1} \frac{r}{p} - [s - (p - 1)]_+.$$

It turns out that, for any given $\beta_0 > 0$ and all $n \geq 1$, by (78),

$$\begin{aligned} \beta_n &= \beta_0 \left(\frac{r}{p} \right)^n - [s - (p - 1)]_+ \sum_{k=0}^{n-1} \left(\frac{r}{p} \right)^k \\ &= \left(\frac{r}{p} \right)^n \left[\beta_0 - [s - (p - 1)]_+ \sum_{j=1}^n \left(\frac{p}{r} \right)^j \right] \\ &= \left(\frac{r}{p} \right)^n \left\{ \beta_0 - [s - (p - 1)]_+ \frac{p}{r - p} \right\} + [s - (p - 1)]_+ \frac{p}{r - p}. \end{aligned}$$

Moreover we have that for all $s \geq 0$,

$$\beta_n \left(\frac{p}{r} \right)^n \rightarrow \beta_0 - [s - (p - 1)]_+ \frac{p}{r - p}, \quad \text{as } n \rightarrow +\infty.$$

Requiring that $\beta_0 > p[s - (p - 1)]_+ / (r - p)$, which will be assumed from now on, then implies that $\beta_n \rightarrow +\infty$ as $n \rightarrow +\infty$. We shall also require $\beta_n \neq (p - 1)$ for any n .

We will explicitly choose a decreasing sequence of radii $0 < R_\infty < \dots < R_n < R_{n-1} < \dots < R_0$ in the next step, in order to estimate the constants. The first iteration step (with $\rho_0 = R_n$, $\rho_1 = R_{n-1}$ and $\beta = \beta_n$) reads:

$$(36) \quad \|u\|_{\beta_n \frac{r}{p}, R_n} \leq I(\beta_n, s, p, R_n, R_{n-1})^{\frac{1}{\beta_n}} \left[\int_{B_{R_{n-1}}} u^{\beta_n + [s - (p-1)]_+} dx \right]^{\frac{1}{\beta_n}} \\ = I_n^{\frac{1}{\beta_n}} \|u\|_{\beta_n \frac{r}{p}, R_{n-1}}^{\frac{r \beta_n - 1}{\beta_n}}$$

where the constants $I_n := I(\beta_n, s, p, R_n, R_{n-1})$ are as (34), that is

$$I_n = \frac{S_p^p}{(R_{n-1} - R_n)^p} \frac{|B_{R_{n-1}}|}{\int_{B_{R_{n-1}}} u^{[s-(p-1)]_+} dx} \left\{ \frac{\beta_n^p A_{s,n}}{|\beta_n - (p-1)|} \right. \\ \left. \times \left(\frac{R_{n-1} - R_n}{R_{n-1}} \right)^p + \left(\frac{R_{n-1} - R_n}{R_n} \right)^p + \frac{\beta_n^p 2^{p-1}}{|\beta_n - (p-1)|^p} \right\},$$

with

$$A_{s,n} := \begin{cases} \frac{\lambda}{p^{p-1}} R_{n-1}^p, & \text{if } s = p - 1, \\ 2^{p-1+d}, & \text{if } s \neq p - 1. \end{cases}$$

Iterating (36), we get

$$\|u\|_{\beta_n \frac{r}{p}, R_n} \leq \dots \leq I_n^{\frac{1}{\beta_n}} I_{n-1}^{\frac{r}{\beta_n}} \dots I_1^{\left(\frac{r}{p}\right)^{n-1} \frac{1}{\beta_n}} \|u\|_{\frac{r}{p} \beta_0, R_0}^{\left(\frac{r}{p}\right)^n \frac{\beta_0}{\beta_n}} \\ = \prod_{j=1}^n I_j^{\left(\frac{r}{p}\right)^{n-j} \frac{1}{\beta_n}} \|u\|_{\frac{r}{p} \beta_0, R_0}^{\left(\frac{r}{p}\right)^n \frac{\beta_0}{\beta_n}},$$

with

$$\beta_0 > [s - (p - 1)]_+ \frac{p}{r - p} \quad \text{or} \quad q = \frac{r}{p} \beta_0 > \bar{q} = [s - (p - 1)]_+ \frac{r}{r - p}.$$

Taking the limit as $n \rightarrow +\infty$ we obtain

$$\|u\|_{\infty, R_\infty} = \lim_{n \rightarrow \infty} \|u\|_{\beta_n \frac{r}{p}, R_n} \leq \lim_{n \rightarrow \infty} \prod_{j=1}^n I_j^{\left(\frac{r}{p}\right)^{n-j} \frac{1}{\beta_n}} \|u\|_{\frac{r}{p} \beta_0, R_0}^{\left(\frac{r}{p}\right)^n \frac{\beta_0}{\beta_n}} \\ = I_\infty \|u\|_{q, R_0}^{\frac{(r-p)q}{(r-p)q - r[s-(p-1)]_+}}.$$

Notice that the last step follows because we shall see below that

$$\prod_{j=1}^n I_j^{\left(\frac{r}{p}\right)^{n-j} \frac{1}{\beta_n}}$$

has a limit I_∞ as $n \rightarrow \infty$. As a consequence of the above estimate we obtain the boundedness of the solution u so that the previous bound holds for any $q > \bar{q}$, as stated.

• STEP 5. *Estimating all the constants.*

Estimating I_∞ . If we are able to prove that for any $j = 1, 2, \dots$

$$(37) \quad I_j \leq I_0 \left(\frac{r}{p} \right)^{pj},$$

holds for some $I_0 > 0$, then we get that

$$\begin{aligned} I_\infty &= \lim_{n \rightarrow \infty} \prod_{j=1}^n I_j^{\left(\frac{r}{p}\right)^{n-j} \frac{1}{\beta_n}} = \lim_{n \rightarrow \infty} \exp \left[\left(\frac{r}{p}\right)^n \frac{1}{\beta_n} \sum_{j=1}^n \left(\frac{p}{r}\right)^j \log(I_j) \right] \\ &\leq \lim_{n \rightarrow \infty} \exp \left\{ \left(\frac{r}{p}\right)^n \frac{1}{\beta_n} \sum_{j=1}^n \left(\frac{p}{r}\right)^j \left[\log I_0 + pj \log \frac{r}{p} \right] \right\} \\ &= \left[I_0 \cdot \left(\frac{r}{p}\right)^{\frac{pr}{r-p}} \right]^{\frac{p}{\beta_0(r-p) - [s-(p-1)]_+ p}} \end{aligned}$$

where we have used the identities (76) and (77).

Estimating I_j . Now we have to prove (37) and so an explicit estimate for I_0 in order to finally obtain (26). Let us choose a decreasing sequence of radii $0 < R_\infty < \dots < R_j < R_{j-1} < \dots < R_0$ such that

$$(R_{j-1} - R_j)^p = (R_0 - R_\infty)^p \frac{c_0^p}{\beta_j^p}, \quad \text{with } c_0 = \left[\sum_{j=1}^\infty \frac{1}{\beta_j} \right]^{-1} < \infty.$$

So

$$\begin{aligned} (38) \quad I_j &\leq \frac{S_p^p \beta_j^p}{c_0^p (R_0 - R_\infty)^p} \frac{|B_{R_0}|}{\int_{B_{R_\infty}} u^{[s-(p-1)]_+} dx} \left\{ \frac{A_{s,0} c_0^p}{|\beta_j - (p-1)|} \left(\frac{R_0 - R_\infty}{R_\infty}\right)^p \right. \\ &\quad \left. + \frac{c_0^p}{\beta_j^p} \left(\frac{R_0 - R_\infty}{R_\infty}\right)^p + \frac{\beta_j^p 2^{p-1}}{|\beta_j - (p-1)|^p} \right\} \end{aligned}$$

recalling that

$$A_{s,0} := \begin{cases} \frac{\lambda}{p^{p-1}} R_0^p, & \text{if } s = p - 1, \\ 2^{p-1+d}, & \text{if } s \neq p - 1. \end{cases}$$

Notice that if we consider the real valued function of a real variable, that is

$$h(t) := \frac{|t - (p-1)|}{t^{p+1}}, \quad \forall t \geq \beta_0,$$

it is easy to show that h is decreasing in $[\beta_0, p-1] \cup [(p^2-1)/p, +\infty)$ and it is increasing otherwise, moreover $\lim_{t \rightarrow \infty} h(t) = 0$. Hence $h(t) \leq \max\{h(\beta_0), h((p^2-1)/p)\}$

for any $t \geq \beta_0$ and so

$$\frac{|\beta_j - (p - 1)|}{\beta_j^{p+1}} \leq \max \left\{ \frac{|\beta_0 - (p - 1)|}{\beta_0^{p+1}}, \left(\frac{p}{p - 1} \right)^p \frac{1}{(p + 1)^{p+1}} \right\} =: c_2.$$

Moreover

$$\beta_j = \left(\frac{r}{p} \right)^j \left\{ \beta_0 - [s - (p - 1)]_+ \frac{p}{r - p} \right\} + [s - (p - 1)]_+ \frac{p}{r - p} \leq \beta_0 \left(\frac{r}{p} \right)^j, \quad \forall j,$$

and

$$(39) \quad \frac{\beta_j}{|\beta_j - (p - 1)|} \leq c_1 := \begin{cases} \frac{\beta_0}{\beta_0 - (p - 1)}, & \text{if } \beta_0 > p - 1, \\ \max_{i=0,1} \frac{\beta_{j_0+i}}{|\beta_{j_0+i} - (p - 1)|}, & \text{if } 0 < \beta_0 < p - 1. \end{cases}$$

As a matter of fact, when $\beta_0 > p - 1$, we have

$$\frac{\beta_j}{|\beta_j - (p - 1)|} \leq \frac{\beta_0}{\beta_0 - (p - 1)}$$

since the one-variable real function

$$\frac{t}{|t - (p - 1)|} = \frac{t}{t - (p - 1)} \quad \text{is decreasing for } t \geq \beta_0 > p - 1.$$

The case $0 < \beta_0 < p - 1$ deserves a further explanation. We define j_0 to be the greatest integer for which $\beta_{j_0} < p - 1$, so that $\beta_{j_0+1} > p - 1$, that is

$$\beta_{j_0} < p - 1 < \beta_{j_0+1} \quad \text{if and only if} \quad j_0 = i.p. \left[\frac{\log \left(\frac{(r-p)(p-1) - p[s-(p-1)]_+}{\beta_0(r-p) - p[s-(p-1)]_+} \right)}{\log \frac{r}{p}} \right]$$

and we shall take $\beta_0 \in (0, p - 1)$ such that

$$(40) \quad \frac{\log \left(\frac{(r-p)(p-1) - p[s-(p-1)]_+}{\beta_0(r-p) - p[s-(p-1)]_+} \right)}{\log \frac{r}{p}} \quad \text{is not an integer.}$$

Summing up, when we consider $0 < \beta_0 < p - 1$, we have to be careful to choose it so that $\beta_j \neq p - 1$ for all j which amounts (40), then we can assure that $\beta_{j_0} < p - 1 < \beta_{j_0+1}$ and we can estimate

$$\frac{\beta_j}{|\beta_j - (p - 1)|} \leq \max_{i=0,1} \frac{\beta_{j_0+i}}{|\beta_{j_0+i} - (p - 1)|} = \max_{i=0,1} \frac{\left(\frac{r}{p} \right)^{j_0+i} \beta_0}{\left(\frac{r}{p} \right)^{j_0+i} \beta_0 - (p - 1)},$$

and (39) is proved. Hence, coming back to (38), we get

$$\begin{aligned}
 I_j &\leq \frac{S_p^p \beta_j^p}{(R_0 - R_\infty)^p} \frac{|B_{R_0}|}{\int_{B_{R_\infty}} u^{[s-(p-1)]_+} dx} \frac{\beta_j}{|\beta_j - (p-1)|} \\
 &\quad \times \left\{ \frac{A_{s,0}}{\beta_j} \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{|\beta_j - (p-1)|}{\beta_j^{p+1}} \right. \\
 &\quad \left. + \left(\frac{\beta_j}{|\beta_j - (p-1)|} \right)^{p-1} \frac{2^{p-1}}{c_0^p} \right\} \\
 &\leq \frac{S_p^p \beta_0^p c_1}{(R_0 - R_\infty)^p} \frac{|B_{R_0}|}{\int_{B_{R_\infty}} u^{[s-(p-1)]_+} dx} \left\{ \frac{A_{s,0}}{\beta_0} \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \right. \\
 &\quad \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p c_2 + \frac{c_1^{p-1} 2^{p-1}}{c_0^p} \right\} \left(\frac{r}{p} \right)^{jp} \\
 &= I_0 \left(\frac{r}{p} \right)^{jp}.
 \end{aligned}$$

Estimate (37) is now proved.

After some simple calculations, the proof is concluded by letting $\beta_0 = pq/r$. □

4.2 - Local upper bounds II. The case of unbounded coefficients.

In this section we establish upper bounds for nonnegative solution to $-\Delta_p u = b(x)u^{p-1}$ on B_R with $b \in L^m(B_R)$ eventually unbounded. These estimates follow from the energy estimates together with the Reverse Poincaré inequalities, which are consequence of Sobolev inequality on balls, see Section 3.

Theorem 4.2 (The Moser iteration). *Let u be a nonnegative weak (sub) solution to $-\Delta_p u = b u^{p-1}$ in B_R , with $b \in L^m(B_R)$, $m > r/(r-p)$ and r as in (14). Then the following bound holds true for any $R_\infty < R_0 < R$ and $q > p-1$*

$$(41) \quad \|u\|_{\infty, R_\infty} \leq \frac{I_{\infty, q}(b)}{(R_0 - R_\infty)^{\frac{pq}{q(r-p)}}} \|u\|_{q, R_0},$$

with constant

$$\begin{aligned}
 I_{\infty,q}(b) &= \left(\frac{S_p^p q^{\frac{mr(p-1)}{m(r-p)-r}}}{c_3^p} \right)^{\frac{r}{q(r-p)}} \left(\frac{r}{p} \right)^{\frac{m^2(p-1)(r+p)}{q(r-p)^2[m(r-p)-r]}} \left[3 \left(\frac{q}{q-(p-1)} \right)^p \right. \\
 &\quad \times \frac{p^p}{q^{\frac{rm(p-1)}{m(r-p)-r}}} + K_{m,p,r} S_p^{\frac{p(mp+r)}{m(r-p)-r}} \left(\frac{q}{q-(p-1)} \right)^{\frac{mr}{m(r-p)-r}} \\
 &\quad \times \frac{2^{\frac{mr+(mp+r)(p-1)}{m(r-p)-r}}}{p^{\frac{rm(p-1)}{m(r-p)-r}}} (R_0 - R_\infty)^p |B_{R_0}|^{\frac{p}{r}} \|b\|_{m,R_0}^{\frac{mr}{m(r-p)-r}} \\
 &\quad \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{1}{(p-1)^{\frac{m(r-p-1)}{m(r-p)-r}}} \right]^{\frac{r}{q(r-p)}} \\
 (42) \quad c_3 &:= \left(\frac{r}{p} \right)^{\frac{mr(p-1)}{p[m(r-p)-r]}} - 1
 \end{aligned}$$

and $K_{m,p,r}$ as in (17).

Remark 4.1. Notice that in the case of bounded coefficients, i. e. $b \in L^\infty(B_R)$, we can pass to the limit as $m \rightarrow \infty$ in the above expression $I_{\infty,q}(b)$ to get

$$\begin{aligned}
 I_{\infty,q}(b) &= \left(S_p^p q^{\frac{r(p-1)}{r-p}} \right)^{\frac{r}{q(r-p)}} \left[\left(\frac{r}{p} \right)^{\frac{r(p-1)}{r-p}} - 1 \right]^{-\frac{r}{q(r-p)}} \left(\frac{r}{p} \right)^{\frac{r^2(p-1)(r+p)}{q(r-p)^3}} \\
 &\quad \times \left[3 \left(\frac{q}{q-(p-1)} \right)^p \frac{p^p}{q^{\frac{r(p-1)}{r-p}}} + S_p^{\frac{p^2}{r-p}} \frac{r-p}{r} \left(\frac{p}{r} \right)^{\frac{p}{r-p}} 2^{\frac{r+p(p-1)}{r-p}} \frac{r}{p^{\frac{r(p-1)}{r-p}}} \right. \\
 &\quad \times \left(\frac{q}{q-(p-1)} \right)^{\frac{r}{r-p}} (R_0 - R_\infty)^p |B_{R_0}|^{\frac{p}{r}} \|b\|_{\infty,R_0}^{\frac{r}{r-p}} \\
 &\quad \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{1}{(p-1)^{\frac{r(p-1)}{r-p}}} \right]^{\frac{r}{q(r-p)}}.
 \end{aligned}$$

Proof. We underline, as in Remark 3.1, that the requirement u belonging to $L^{\alpha+(p-1)}(B_\rho)$ needed at each iteration step in order to apply the reverse Poincaré inequalities of Theorem 3.1, will be dispensed by the previous one. Therefore, we only have to ensure the integrability condition at the first step, which gives rise to condition $m > r/(r-p)$.

• **STEP 1. Sobolev and Reverse Poincaré inequalities.** We start considering the radii $R_\infty < \rho_1 < \rho_0 < R_0$ and we use (13) on the ball B_{ρ_1} with $g = u^{[\alpha+(p-1)]/p}$, for some $\alpha > 0$, to get

$$\left[\int_{B_{\rho_1}} u^{\frac{\alpha+(p-1)r}{p}} dx \right]^{\frac{p}{r}} \leq S_p^p \left[\int_{B_{\rho_1}} |\nabla(u^{\frac{\alpha+(p-1)}{p}})|^p dx + \frac{1}{\rho_1^p} \int_{B_{\rho_1}} u^{\alpha+(p-1)} dx \right].$$

To estimate the first term in the right hand side of the previous inequality we use Theorem 3.1 (see Remark 3.2), we get

$$(43) \quad \left[\int_{B_{\rho_1}} u^{\frac{\alpha+(p-1)r}{p}} dx \right]^{\frac{p}{r}} \leq S_p^p \left[A(b) + \frac{1}{\rho_1^p} \right] \int_{B_{\rho_0}} u^{\alpha+(p-1)} dx$$

with $A(b)$ as in (22) and $K_{m,p,r}$ in (17).

Notice that

$$\begin{aligned} A(b) &\leq \frac{[\alpha + (p-1)]^{\frac{mr(p-1)}{m(r-p)-r}}}{(\rho_0 - \rho_1)^p} \left\{ 3 \left(\frac{\alpha + (p-1)}{\alpha} \right)^p \frac{p^p}{[\alpha + (p-1)]^{\frac{mr(p-1)}{m(r-p)-r}}} \right. \\ &\quad + S_p^{\frac{p(mp+r)}{m(r-p)-r}} K_{m,p,r} \frac{2^{\frac{mr+(mp+r)(p-1)}{m(r-p)-r}}}{p^{\frac{mr(p-1)}{m(r-p)-r}}} \left(\frac{\alpha + (p-1)}{\alpha} \right)^{\frac{mr}{m(r-p)-r}} \\ &\quad \left. \times (\rho_0 - \rho_1)^p |B_{R_0}|^{\frac{p}{r}} \|b\|_{m,R_0}^{\frac{rm}{m(r-p)-r}} \right\}. \end{aligned}$$

Hence, we get, since $\alpha > 0$,

$$\begin{aligned} S_p^p \left[A(b) + \frac{1}{\rho_1^p} \right] &\leq \frac{S_p^p [\alpha + (p-1)]^{\frac{mr(p-1)}{m(r-p)-r}}}{(\rho_0 - \rho_1)^p} \left\{ 3 \left(\frac{\alpha + (p-1)}{\alpha} \right)^p \right. \\ &\quad \times \frac{p^p}{[\alpha + (p-1)]^{\frac{mr(p-1)}{m(r-p)-r}}} + S_p^{\frac{p(mp+r)}{m(r-p)-r}} K_{m,p,r} \\ &\quad \times \frac{2^{\frac{mr+(mp+r)(p-1)}{m(r-p)-r}}}{p^{\frac{mr(p-1)}{m(r-p)-r}}} \left(\frac{\alpha + (p-1)}{\alpha} \right)^{\frac{mr}{m(r-p)-r}} \\ &\quad \times (R_0 - R_\infty)^p |B_{R_0}|^{\frac{p}{r}} \|b\|_{m,R_0}^{\frac{rm}{m(r-p)-r}} \\ &\quad \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{1}{(p-1)^{\frac{mr(p-1)}{m(r-p)-r}}} \right\}. \end{aligned}$$

• **STEP 2. The Moser iteration.** We now fix $\beta_0 = \alpha + (p - 1) > p - 1$ and we define the sequence

$$\beta_n = \frac{r}{p} \beta_{n-1} = \left(\frac{r}{p}\right)^n \beta_0$$

and that of radii $R_\infty < \dots < R_n < R_{n-1} < \dots < R_0$ such that

$$(44) \quad (R_{n-1} - R_n)^p = c_3^p (R_0 - R_\infty)^p \left(\frac{p}{r}\right)^{\frac{mr(p-1)n}{m(r-p)-r}}$$

with

$$c_3 := \left[\sum_{k=1}^{\infty} \left(\frac{p}{r}\right)^{\frac{mr(p-1)k}{p(m(r-p)-r)}} \right]^{-1} = \left(\frac{r}{p}\right)^{\frac{mr(p-1)}{p(m(r-p)-r)}} - 1,$$

recalling (76), so that

$$\sum_{n=1}^{\infty} (R_{n-1} - R_n) = (R_0 - R_\infty).$$

With these choices inequality (43), in which $\alpha + (p - 1)$ is replaced by $\beta_{n-1} > p - 1$ and ρ_1, ρ_0 by R_n, R_{n-1} respectively, reads

$$\begin{aligned} \left[\int_{B_{R_n}} w^{\beta_n} dx \right]^{\frac{p}{r}} &\leq \frac{S_p^p \beta_{n-1}^{\frac{mr(p-1)}{m(r-p)-r}}}{(R_{n-1} - R_n)^p} \left\{ \left(\frac{\beta_{n-1}}{\beta_{n-1} - (p-1)} \right)^p \frac{p^p}{\beta_{n-1}^{\frac{mr(p-1)}{m(r-p)-r}}} \right. \\ &\quad + S_p^{\frac{p(mp+r)}{m(r-p)-r}} \frac{2^{\frac{mr+(mp+r)(p-1)}{m(r-p)-r}}}{p^{\frac{mr(p-1)}{m(r-p)-r}}} \left(\frac{\beta_{n-1}}{\beta_{n-1} - (p-1)} \right)^{\frac{mr}{m(r-p)-r}} \\ &\quad \times K_{m,p,r} (R_0 - R_\infty)^p |B_{R_0}|^{\frac{p}{r}} \|b\|_{m,R_0}^{\frac{mr}{m(r-p)-r}} \\ &\quad \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{1}{(p-1)^{\frac{mr(p-1)}{m(r-p)-r}}} \right\} \int_{B_{R_{n-1}}} w^{\beta_{n-1}} dx \\ &= I_{n-1} \int_{B_{R_{n-1}}} w^{\beta_{n-1}} dx. \end{aligned}$$

Letting $Y_n := \|u\|_{\beta_n, R_n}$, we have obtained

$$Y_n \leq I_{n-1}^{\left(\frac{p}{r}\right)^{n-1}} \frac{1}{\beta_0} Y_{n-1}.$$

If we prove $I_{n-1} \leq C^{n-1} I_0$, for some $I_0, C > 0$, we can apply Lemma 8.3 with $\sigma = 1/\beta_0$ and $\theta = p/r \in (0, 1)$ to get

$$Y_\infty \leq I_0^{\frac{\sigma}{1-\theta}} C^{\frac{\sigma\theta}{(1-\theta)^2}} Y_0 \quad \text{which is} \quad \|u\|_{\infty, R_\infty} \leq I_{\infty, \beta_0}(b) \|u\|_{\beta_0, R_0}$$

with

$$\begin{aligned}
 I_\infty(b) = & \left\{ \frac{S_p^p \beta_0^{\frac{mr(p-1)}{m(r-p)-r}}}{c_3^p (R_0 - R_\infty)^p} \left[3 \left(\frac{\beta_0}{\beta_0 - (p-1)} \right)^p \frac{p^p}{\beta_0^{\frac{mr(p-1)}{m(r-p)-r}}} \right. \right. \\
 & + \frac{2^{\frac{mr+(mp+r)(p-1)}{m(r-p)-r}}}{p^{\frac{mr(p-1)}{m(r-p)-r}}} K_{m,p,r} \left(\frac{\beta_0}{\beta_0 - (p-1)} \right)^{\frac{mr}{m(r-p)-r}} \\
 & \times (R_0 - R_\infty)^p |B_{R_0}|^{\frac{p}{r}} \|b\|_{m,R_0}^{\frac{mr}{m(r-p)-r}} \\
 & \left. \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{1}{(p-1)^{\frac{mr(p-1)}{m(r-p)-r}}} \right] \right\} \frac{r}{\beta_0^{\frac{r}{r-p}}} \left(\frac{r}{p} \right)^{\frac{mr^2(p-1)(r+p)}{\beta_0^{lm(r-p)-r}(r-p)^2}}.
 \end{aligned}$$

The proof is concluded once we show that $I_{n-1} \leq C^{n-1} I_0$, for some $I_0, C > 0$ and let $\beta_0 = q > p - 1$. To prove the above estimate for I_{n-1} , it is sufficient to note that

$$\begin{aligned}
 I_{n-1} \leq & \frac{S_p^p \beta_0^{\frac{mr(p-1)}{m(r-p)-r}}}{c_3^p (R_0 - R_\infty)^p} \left\{ 3 \left(\frac{\beta_0}{\beta_0 - (p-1)} \right)^p \frac{p^p}{\beta_0^{\frac{mr(p-1)}{m(r-p)-r}}} + \frac{2^{\frac{mr+(mp+r)(p-1)}{m(r-p)-r}}}{p^{\frac{mr(p-1)}{m(r-p)-r}}} \right. \\
 & \times S_p^{\frac{p(mp+r)}{m(r-p)-r}} K_{m,p,r} \left(\frac{\beta_0}{\beta_0 - (p-1)} \right)^{\frac{mr}{m(r-p)-r}} \\
 & \times (R_0 - R_\infty)^p |B_{R_0}|^{\frac{p}{r}} \|b\|_{m,R_0}^{\frac{mr}{m(r-p)-r}} \\
 & \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{1}{(p-1)^{\frac{mr(p-1)}{m(r-p)-r}}} \right\} \left(\frac{r}{p} \right)^{\frac{mr(p-1)}{m(r-p)-r}} \left(\frac{r}{p} \right)^{\frac{2mr(p-1)(n-1)}{m(r-p)-r}} \\
 =: & I_0 C^{n-1},
 \end{aligned}$$

where we have used (44), the definition of β_{n-1} and the following facts

$$\frac{\beta_{n-1}}{\beta_{n-1} - (p-1)} \leq \frac{\beta_0}{\beta_0 - (p-1)}, \text{ for } \beta_0 > p - 1 \text{ and } \frac{1}{\beta_{n-1}} \leq \frac{1}{\beta_0}.$$

□

Theorem 4.3 (Extending Local Upper Bounds). *Let u be a nonnegative weak solution to $-\Delta_p u = b u^{p-1}$ in B_R , with $p > 1$, $b \in L^m(B_R)$, $m > r/(r-p)$ and r as in (14). Then the following bound holds for any $B_{R_\infty} \subset B_{R_0} \subseteq B_R$ and for any $q > 0$*

$$\|u\|_{\infty, R_\infty} \leq \frac{A_q^{(1)}}{(R_0 - R_\infty)^{\frac{pr}{q(r-p)}}} \left[A_q^{(2)} + A_q^{(3)} \|b\|_{m,R_0}^{\frac{mr}{m(r-p)-r}} \right]^{\frac{r}{q(r-p)}} \|u\|_{q,R_0}$$

the constants $A_q^{(i)}$, for $i = 1, 2, 3$ depend on $d, p, s, q, r, R_0, R_\infty$, see an explicit expression in formulas (45) below.

Remark on the constant. The proof below allows us to find an explicit expression of the constant:

$$(45) \quad A_q^{(1)} := \left(\frac{S_p^p q^{\frac{mr(p-1)}{m(r-p)-r}}}{c_3^p} \right)^{\frac{r}{q(r-p)}} \left(\frac{r}{p} \right)^{\frac{m^2(p-1)(r+p)}{q(r-p)^2[m(r-p)-r]}}$$

if $q > p - 1$,

$$A_q^{(1)} := \mathfrak{z} \left(\frac{S_p^p [q + (p-1)]^{\frac{mr(p-1)}{m(r-p)-r}}}{c_3^p} \right)^{\frac{r}{q(r-p)}} \left(\frac{r}{p} \right)^{\frac{m^2(p-1)(r+p)}{q(r-p)^2[m(r-p)-r]}} \\ \times \left(\frac{pr}{q(r-p)} \right)^{\frac{pr}{q(r-p)}} 2^{\frac{p-1}{q} + \frac{2pr}{q(r-p)}}$$

if $0 < q \leq p - 1$,

$$A_q^{(2)} := \mathfrak{z} \left(\frac{q}{q - (p-1)} \right)^p \frac{p^p}{q^{\frac{mr(p-1)}{m(r-p)-r}}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{1}{(p-1)^{\frac{mr(p-1)}{m(r-p)-r}}}$$

if $q > p - 1$,

$$A_q^{(2)} := \left(\frac{q + (p-1)}{q} \right)^p \frac{p^p}{[q + (p-1)]^{\frac{mr(p-1)}{m(r-p)-r}}} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \frac{\mathfrak{z}}{(p-1)^{\frac{mr(p-1)}{m(r-p)-r}}}$$

if $0 < q \leq p - 1$ and

$$A_q^{(3)} := \frac{2^{\frac{mr+(mp+r)(p-1)}{m(r-p)-r}} K_{m,p,r}}{p^{\frac{mr(p-1)}{m(r-p)-r}}} \left(\frac{q S_p^{\frac{p(mp+r)}{mr}}}{q - (p-1)} \right)^{\frac{mr}{m(r-p)-r}} (R_0 - R_\infty)^p |B_{R_0}|^{\frac{p}{r}}$$

if $q > p - 1$

$$A_q^{(3)} := K_{m,p,r} \left(\frac{[q + (p-1)] S_p^{\frac{p(mp+r)}{mr}}}{q p^{p-1} 2^{\frac{mr+(mp+r)(p-1)}{mr}}} \right)^{\frac{mr}{m(r-p)-r}} (R_0 - R_\infty)^p |B_{R_0}|^{\frac{p}{r}}$$

if $0 < q \leq p - 1$, with c_3 and $K_{m,p,r}$, respectively as in (42) and (17).

Proof. The statement of the theorem, in the case $q > p - 1$, easily follows from Theorem 4.2. When $0 < q \leq p - 1$ we apply Lemma 8.5. Indeed by Theorem 4.2 (with q substitutes for $q + (p - 1) > p - 1$), we have

$$\|u\|_{\infty, R_\infty} \leq \frac{I_{\infty, q+(p-1)}(b)}{(R_0 - R_\infty)^{\frac{pr}{q+(p-1)(r-p)}}} \|u\|_{q+(p-1), R_0}$$

and hence we arrive at the desired result using Lemma 8.5 with $q_2 = \infty$, $q_1 = q + (p - 1)$, $K = I_{\infty, q+(p-1)}(b)$ and $\gamma = pr/\{[q + (p - 1)](r - p)\}$. \square

The above theorem has the following important consequence, when applied to the equation $-\Delta_p u = \lambda u^s$.

Theorem 4.4 (Local Upper Bounds, second form). *Let u be a nonnegative weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $p > 1$, $\lambda > 0$, $s > p - 1$ and r as in (14). If $u \in L^{\bar{m}}(\Omega)$, with $\bar{m} > r[s - (p - 1)]/(r - p)$, then the following bound holds for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$ and for any $q > 0$*

$$\begin{aligned} \|u\|_{\infty, R_\infty} &\leq \frac{A_q^{(1)} \left[A_q^{(2)} + A_q^{(3)} \lambda^{\frac{\bar{m}r}{\bar{m}(r-p)-r[s-(p-1)]}} \|u\|_{\frac{\bar{m}}{m}, R_0}^{\frac{\bar{m}r[s-(p-1)]}{\bar{m}(r-p)-r[s-(p-1)]}} \right]^{\frac{r}{q(r-p)}}}{(R_0 - R_\infty)^{\frac{pr}{q(r-p)}}} \\ &\quad \times \|u\|_{q, R_0}, \end{aligned}$$

where $A_q^{(1)}$, $A_q^{(2)}$ and $A_q^{(3)}$ are as in Theorem 4.3.

Proof. Since u is a solution to $-\Delta_p u = \lambda u^s$ in B_R , then u is also a solution to $-\Delta_p u = b u^{p-1}$ on B_R with $b = \lambda u^{s-(p-1)}$. Therefore we need to assume $u^{s-(p-1)} \in L^m(B_R)$, with $m > r/(r - p)$ that it is equivalent to require $u \in L^{\bar{m}}(B_R)$, with $\bar{m} = m[s - (p - 1)] > r[s - (p - 1)]/(r - p)$. So that

$$\begin{aligned} \|b\|_{\frac{mr}{m}, R_0}^{\frac{mr}{\bar{m}(r-p)-r}} &= \left[\lambda^m \int_{B_{R_0}} u^{m[s-(p-1)]} dx \right]^{\frac{r}{\bar{m}(r-p)-r}} \\ &= \lambda^{\frac{\bar{m}r}{\bar{m}(r-p)-r[s-(p-1)]}} \|u\|_{\frac{\bar{m}}{m}, R_0}^{\frac{\bar{m}r[s-(p-1)]}{\bar{m}(r-p)-r[s-(p-1)]}}. \end{aligned}$$

Hence the result follows from Theorem 4.3. \square

5 - Local lower bounds via Moser iteration

In this section we prove quantitative local lower bounds for nonnegative weak solutions to $-\Delta_p u = \lambda u^s$. The strategy to prove the lower bounds is classical, and combines a lower Moser iteration with some reverse Hölder inequalities obtained via a John-Nirenberg type Lemma. Since we are interested in keeping track of all (the

relevant) constants, we need a quantitative version of a John-Nirenberg type Lemma to obtain quantitative reverse Hölder inequalities; this has been done in [4] and the proofs of [4] also adapt to the current setting with minor modifications that we give in Appendix 8.1.

We first show how the lower Moser iteration proves quantitative local lower bounds in a general form, that hold in the whole range $0 \leq s < r - 1$. In the next subsection, we will improve the results in a smaller range, namely $p - 1 < s < s_c^* = r(p - 1)/p$.

Theorem 5.1 (Local Lower Estimates). *Let $\Omega \subset \mathbb{R}^d$. Let u be a nonnegative local weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $p > 1$, $\lambda \geq 0$ and $0 \leq s < s_c = r - 1$, r as in (14). Then for any $\varepsilon > 0$, for any*

$$0 < q \leq \frac{(p - 1)^{\frac{2}{p}} 2^{\frac{(d-1)(p-1)}{p}}}{p \omega_d^{\frac{2}{d}} d [e(d - 1) + \varepsilon]} =: \underline{q}_\varepsilon$$

and for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$ the following bound holds

$$\inf_{x \in B_{R_\infty}} u(x) = \|u\|_{-\infty, R_\infty} \geq I_{-\infty, q} \frac{\|u\|_{q, R_0}}{|B_{R_0}|^{\frac{1}{q}}}$$

where

$$(46) \quad I_{-\infty, q} := \left[\frac{(R_0 - R_\infty) R_\infty}{R_0^{\frac{d(r-p)}{rp}}} \right]^{\frac{pr}{q(r-p)}} \left[\frac{1}{S_p^p [2^{p-1} R_\infty^p + (R_0 - R_\infty)^p] 2^{\frac{pr}{r-p}}} \right]^{\frac{r}{q(r-p)}} \times \left[\frac{\varepsilon}{2^d (e d + \varepsilon) \sqrt{\omega_d}} \right]^{\frac{2}{q}}.$$

Proof. The proof is divided in two steps.

• **STEP 1.** In this step we want to prove $L^{-q} - L^{-\infty}$ local estimates via Moser iteration. Consider $\alpha < -(p - 1)$, choosing ϕ as in Lemma 2.2 in the estimate (4), we obtain

$$\int_{B_{\rho_1}} |\nabla[(u + \delta)^{\frac{\alpha+p-1}{p}}]|^p dx \leq \frac{2^{p-1} |\alpha + (p - 1)|^p}{|\alpha|^p (\rho_0 - \rho_1)^p} \int_{B_{\rho_0}} (u + \delta)^{\alpha+p-1} dx.$$

Applying now the Sobolev inequality (13) on the ball B_{ρ_1} , let ρ_1 be any real positive number such that $B_{\rho_1} \subset \subset \Omega$, one gets

$$\begin{aligned}
 \left[\int_{B_{\rho_1}} (u + \delta)^{\frac{\alpha+p-1}{p}r} dx \right]^{\frac{p}{r}} &\leq S_p^p \left[\int_{B_{\rho_1}} |\nabla[(u + \delta)^{\frac{\alpha+p-1}{p}}]|^p dx \right. \\
 &\quad \left. + \frac{1}{\rho_1^p} \int_{B_{\rho_1}} (u + \delta)^{\alpha+p-1} dx \right] \\
 &\leq S_p^p \left[\frac{2^{p-1} |\alpha + (p-1)|^p}{|\alpha|^p (\rho_0 - \rho_1)^p} + \frac{1}{\rho_1^p} \right] \\
 &\quad \times \int_{B_{\rho_0}} (u + \delta)^{\alpha+p-1} dx \\
 &\leq S_p^p \left[\frac{2^{p-1}}{(\rho_0 - \rho_1)^p} + \frac{1}{\rho_1^p} \right] \int_{B_{\rho_0}} (u + \delta)^{\alpha+p-1} dx,
 \end{aligned}$$

since $|\alpha + (p - 1)|/|\alpha| < 1$ for any $\alpha < -(p - 1)$. Let for a given $\gamma_0 < 0$,

$$\gamma_n = \frac{r}{p} \gamma_{n-1} = \left(\frac{r}{p}\right)^n \gamma_0.$$

Notice that $\gamma_n \rightarrow -\infty$ monotonically. The above inequality, with $\alpha = \alpha_n$, $\gamma_{n-1} = \alpha_n + (p - 1) < 0$ and $\rho_1 = R_n < \rho_0 = R_{n-1}$ reads

$$\begin{aligned}
 \|u + \delta\|_{\gamma_n, R_n} &\geq \left[S_p^p \left(\frac{2^{p-1}}{(R_{n-1} - R_n)^p} + \frac{1}{R_n^p} \right) \right]^{\frac{1}{\gamma_{n-1}}} \\
 &\quad \times \left[\int_{B_{R_{n-1}}} (u + \delta)^{\gamma_{n-1}} dx \right]^{\frac{1}{\gamma_{n-1}}} \\
 &=: I_n^{\frac{1}{\gamma_{n-1}}} \|u + \delta\|_{\gamma_{n-1}, R_{n-1}}.
 \end{aligned}$$

The iteration is simple now, and gives

$$(47) \quad \|u + \delta\|_{\gamma_n, R_n} \geq I_n^{\frac{1}{\gamma_{n-1}}} I_{n-1}^{\frac{1}{\gamma_{n-2}}} \dots I_1^{\frac{1}{\gamma_0}} \|u + \delta\|_{\gamma_0, R_0} = \prod_{j=1}^n I_j^{\frac{1}{\gamma_{j-1}}} \|u + \delta\|_{\gamma_0, R_0},$$

where we have chosen $0 < R_\infty < \dots < R_n < R_{n-1} < \dots < R_0$ such that

$$\sum_{j=1}^\infty (R_{j-1} - R_j) = R_0 - R_\infty \quad \text{and} \quad R_{j-1} - R_j = \frac{R_0 - R_\infty}{2^j}$$

so that

$$I_j = S_p^p \left(\frac{2^{p-1} 2^{pj}}{(R_0 - R_\infty)^p} + \frac{1}{R_j^p} \right) \leq \frac{S_p^p}{(R_0 - R_\infty)^p} \left[2^{p-1} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \right] 2^{pj}$$

$$=: I_0 2^{pj}$$

and

$$\begin{aligned} \prod_{j=1}^n (I_0 2^{pj})^{\frac{1}{j-1}} &= \exp \left\{ \sum_{j=1}^n \frac{1}{\gamma_{j-1}} \log (I_0 2^{pj}) \right\} \\ &= \exp \left\{ \frac{1}{\gamma_0} \frac{r}{p} \sum_{j=1}^n \left(\frac{p}{r} \right)^j \log (I_0 2^{pj}) \right\} \\ &= \exp \left\{ \frac{1}{\gamma_0} \frac{r}{p} \log I_0 \sum_{j=1}^n \left(\frac{p}{r} \right)^j + \frac{1}{\gamma_0} r \log 2 \sum_{j=1}^n j \left(\frac{p}{r} \right)^j \right\} \\ &= I_0^{\frac{1}{\gamma_0} \frac{r}{p}} \sum_{j=1}^n \left(\frac{p}{r} \right)^j 2^{\frac{r}{p} \sum_{j=1}^n j \left(\frac{p}{r} \right)^j} \\ &\quad \downarrow \quad n \rightarrow \infty \\ &= I_0^{\frac{1}{\gamma_0} \frac{r}{p}} \sum_{j=1}^{\infty} \left(\frac{p}{r} \right)^j 2^{\frac{r}{p} \sum_{j=1}^{\infty} j \left(\frac{p}{r} \right)^j}. \end{aligned}$$

Using (76) and (77), we get

$$\prod_{j=1}^n (I_0 2^{pj})^{\frac{1}{j-1}} \xrightarrow{n \rightarrow \infty} I_0^{\frac{1}{\gamma_0} \frac{r}{r-p}} 2^{\frac{r}{\gamma_0} \frac{pr}{(r-p)^2}} = (I_0 2^{\frac{rp}{r-p}})^{\frac{r}{(r-p)}}.$$

We can now take the limit in (47) to get for any $\gamma_0 < 0$

$$(48) \quad \begin{aligned} \|u + \delta\|_{-\infty, R_\infty} &\geq \lim_{n \rightarrow \infty} \|u + \delta\|_{\gamma_n, R_n} \geq \lim_{n \rightarrow \infty} \prod_{j=1}^n (I_0 2^{pj})^{\frac{1}{j-1}} \|u + \delta\|_{\gamma_0, R_0} \\ &= \left\{ \frac{S_p^p}{(R_0 - R_\infty)^p} \left[2^{p-1} + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \right] 2^{\frac{pr}{r-p}} \right\}^{\frac{r}{(r-p)}} \\ &\quad \times \|u + \delta\|_{\gamma_0, R_0}. \end{aligned}$$

• **STEP 2. Reverse Hölder inequalities.** Joining inequality (48) and (75) and letting $\gamma_0 = -q$, for any

$$0 < q \leq \frac{(p-1)^{\frac{2}{p}} 2^{\frac{(d-1)(p-1)}{p}}}{p \omega_d^2 d [e(d-1) + \varepsilon]} = \underline{q}_\varepsilon$$

we obtain

$$\begin{aligned} \|u + \delta\|_{-\infty, R_\infty} &\geq \left[\frac{(R_0 - R_\infty)^p R_\infty^p}{S_p^p [2^{p-1} R_\infty^p + (R_0 - R_\infty)^p] 2^{\frac{rp}{r-p}} R_0^{\frac{d(r-p)}{r}}} \right]^{\frac{r}{q(r-p)}} \\ &\quad \times \left[\frac{\varepsilon}{2^d (e d + \varepsilon) \sqrt{\omega_d}} \right]^{\frac{2}{q}} \frac{\|u + \delta\|_{q, R_0}}{|B_{R_0}|^{\frac{1}{q}}}. \end{aligned}$$

To conclude the proof it is sufficient let $\delta \rightarrow 0^+$. □

5.1 - Reverse Hölder inequalities and additional local lower bounds

In this section we will prove first a more precise quantitative reverse Hölder inequality, that holds in the smaller range of exponents $s > p - 1$. We have in mind to join local upper and lower estimates to get a clean form of Harnack inequality (see next section). The difficulty here is that the lower bound of the previous section has the form of reverse smoothing effect from L^q to $L^{-\infty}$ for a suitable explicit q , which can be very small, sometimes too small: we need to reach higher values of q , namely above $\bar{q} := r[s - (p - 1)]/(r - p)$ and this will be possible through a reverse Hölder inequality, that holds only when $p - 1 < s < r(p - 1)/p = s_c^*$. Under no further assumptions on the solution at hand, it is impossible -to our knowledge- to extend this reverse Hölder inequality to higher values of s in a quantitative way.

Proposition 5.1 (Reverse Hölder inequalities). *Let $\Omega \subset \mathbb{R}^d$ and let $\lambda \geq 0$. Let u be a nonnegative local weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $p - 1 < s < r(p - 1)/p = s_c^*$. Let $B_{\rho_1} \subset B_{\rho_0} \subset \subset \Omega$. Then*

$$(49) \quad \frac{\|u\|_{q_0, \rho_1}}{|B_{\rho_1}|^{\frac{1}{q_0}}} \leq I_{q_0, q} \frac{\|u\|_{q, \rho_0}}{|B_{\rho_0}|^{\frac{1}{q}}},$$

for any $q \in (0, q_0]$, with $\bar{q} < q_0 < s_c^*$,

$$(50) \quad \begin{aligned} I_{q_0, q} &:= \left\{ S_p^p \left[\frac{2^{p-1} p^p q_0^p}{[r(p-1) - pq_0]^p} + \left(\frac{\rho_0 - \rho_1}{\rho_1} \right)^p \right]^{\frac{r}{p q_0}} \right. \\ &\quad \times \left. \left[\frac{\rho_0^{\frac{d(\frac{1}{p} - \frac{1}{r})}}{\rho_0 - \rho_1}}{\omega_d^{\frac{1}{p} - \frac{1}{r}}} \right]^{\frac{r}{q_0}} \left(\frac{\rho_0}{\rho_1} \right)^{\frac{d}{q_0}} \right. \end{aligned}$$

if $pq_0/r \leq q \leq q_0$ and

$$\begin{aligned}
 I_{q_0,q} &:= 3 \cdot 2^{\frac{pq_0-rq}{(r-p)q}} \left\{ S_p^p \left[\frac{2^{p-1} p^p q_0^p}{[r(p-1) - pq_0]^p} \left(\frac{\rho_1}{\rho_0 - \rho_1} \right)^p + 1 \right] \right\}^{\frac{r(q_0-q)}{(r-p)qq_0}} \\
 (51) \quad &\times \left(\frac{4rp(q_0-q)\omega_d^{\frac{r-p}{rp}}}{(r-p)q_0q} \right)^{\frac{rp(q_0-q)}{(r-p)qq_0}} \frac{\rho_0^{\frac{d}{q}}}{\rho_1^{\frac{p^r(q_0-q)}{(r-p)qq_0} + \frac{d}{q_0}}},
 \end{aligned}$$

if $0 < q < pq_0/r$.

Remark 5.1. We note that the interval in which q_0 can vary is not empty, since we are assuming $s < s_c^*$.

Proof. Let $-(p-1) < \alpha < 0$. Consider the energy estimate (6). It implies, using ϕ as in Lemma 2.2,

$$\int_{B_{\rho_1}} |\nabla(u^{\frac{\alpha+(p-1)}{p}})|^p dx \leq \frac{2^{p-1}[\alpha + (p-1)]^p}{|\alpha|^p(\rho_0 - \rho_1)^p} \int_{B_{\rho_0}} u^{\alpha+(p-1)} dx.$$

Applying now Sobolev inequality (13) with $g = u^{[\alpha+(p-1)]/p}$ on the ball B_{ρ_1} we arrive at

$$\begin{aligned}
 \left[\int_{B_{\rho_1}} u^{\frac{\alpha+(p-1)}{p}r} dx \right]^{\frac{p}{r}} &\leq S_p^p \left[\int_{B_{\rho_1}} |\nabla(u^{\frac{\alpha+(p-1)}{p}})|^p dx \right. \\
 &\quad \left. + \frac{1}{\rho_1^{\frac{p}{p}}} \int_{B_{\rho_1}} u^{\alpha+(p-1)} dx \right] \\
 &\leq S_p^p \left[\frac{2^{p-1}[\alpha + (p-1)]^p}{|\alpha|^p(\rho_0 - \rho_1)^p} + \frac{1}{\rho_1^{\frac{p}{p}}} \right] \int_{B_{\rho_0}} u^{\alpha+(p-1)} dx.
 \end{aligned}$$

Letting $0 < \alpha + (p-1) =: \beta < p-1$, we get

$$\begin{aligned}
 (52) \quad \left[\int_{B_{\rho_1}} u^{\beta \frac{r}{p}} dx \right]^{\frac{p}{r}} &\leq \frac{S_p^{\frac{p}{r}}}{(\rho_0 - \rho_1)^{\frac{p}{r}}} \left[\frac{2^{p-1} \beta^p}{|\beta - (p-1)|^p} + \left(\frac{\rho_0 - \rho_1}{\rho_0} \right)^{p \frac{1}{r}} \right]^{\frac{1}{r}} \\
 &\quad \times \left[\int_{B_{\rho_0}} u^{\beta} dx \right]^{\frac{1}{r}}.
 \end{aligned}$$

Let $q = r\beta/p$, then

$$\bar{q} = \frac{r[s - (p - 1)]}{r - p} < q < \frac{r}{p}(p - 1) \quad \text{imply} \quad \frac{p[s - (p - 1)]}{r - p} < \beta < p - 1.$$

We note that the interval in which β can vary is compatible with the request $0 < \beta < p - 1$ and it is not empty since we are assuming $p - 1 < s < s_c^*$. With this choice, from (52) we get, for any $B_{\rho_1} \subset B_{\rho_0} \subset \subset \Omega$,

$$(53) \quad \|u\|_{q,\rho_1} \leq \left\{ S_p^p \left[\frac{2^{p-1} p^p q^p}{[r(p-1) - pq]^p} + \left(\frac{\rho_0 - \rho_1}{\rho_1} \right)^p \right] \right\}^{\frac{r}{pq}} \frac{\|u\|_{\frac{p}{r}q,\rho_0}}{(\rho_0 - \rho_1)^{\frac{r}{q}}}.$$

We consider separately the case $pq/r \leq q_0 \leq q$ and the case $0 < q_0 < pq/r$. In the first case we can use Hölder inequality in (53), to obtain

$$\begin{aligned} \|u\|_{q,\rho_1} &\leq \left\{ S_p^p \left[\frac{2^{p-1} p^p q^p}{[r(p-1) - pq]^p} + \left(\frac{\rho_0 - \rho_1}{\rho_1} \right)^p \right] \right\}^{\frac{r}{pq}} \left[\frac{\rho_0^{\frac{d}{p}} \omega_d^{\frac{1}{p} - \frac{1}{r}}}{(\rho_0 - \rho_1) \rho_1^{\frac{d}{r}}} \right]^{\frac{r}{q}} \\ &\quad \times |B_{\rho_1}|^{\frac{1}{q}} \frac{\|u\|_{q_0,\rho_0}}{|B_{\rho_0}|^{\frac{1}{q_0}}}, \end{aligned}$$

which is (49). On the other hand, when $0 < q_0 < pq/r$, we can use inequality (53) rewritten as

$$\begin{aligned} \|u\|_{q,\rho_1} &\leq \left\{ S_p^p \left[\frac{2^{p-1} p^p q^p}{[r(p-1) - pq]^p} + \left(\frac{\rho_0 - \rho_1}{\rho_1} \right)^p \right] \right\}^{\frac{r}{pq}} \frac{\|u\|_{\frac{p}{r}q,\rho_0}}{(\rho_0 - \rho_1)^{\frac{r}{q}}} \\ &=: \frac{K}{(\rho_0 - \rho_1)^{\frac{r}{q}}} \|u\|_{\frac{p}{r}q,\rho_0}, \end{aligned}$$

so that Lemma 8.5 applied with $q_2 = q$, $q_1 = pq/r$ and $\gamma = r/q$ gives that for all $0 < q_0 < pq/r$

$$\begin{aligned} \|u\|_{q,\rho_1} &\leq 3 \cdot 2^{\frac{pq-rq_0}{q_0(r-p)}} \left\{ S_p^p \left[\frac{2^{p-1} p^p q^p}{[r(p-1) - pq]^p} \left(\frac{\rho_1}{\rho_0 - \rho_1} \right)^p + 1 \right] \right\}^{\frac{r(q-q_0)}{q_0(r-p)}} \\ &\quad \times \left(\frac{4rp(q-q_0)}{q q_0(r-p)} \right)^{\frac{pr(q-q_0)}{q_0 q(r-p)}} \frac{\rho_0^{\frac{d}{q_0}} \omega_d^{\frac{1}{q_0}}}{\rho_1^{\frac{d}{q} + \frac{rp(q-q_0)}{q_0(r-p)}} \omega_d^{\frac{1}{q}}} |B_{\rho_1}|^{\frac{1}{q}} \frac{\|u\|_{q_0,\rho_0}}{|B_{\rho_0}|^{\frac{1}{q_0}}}. \end{aligned}$$

□

As a consequence of the above proposition we can improve the local lower bounds of Theorem 5.1 in this good supercritical range.

Theorem 5.2 (Local Lower Estimates). *Let $\Omega \subset \mathbb{R}^d$ and let $\lambda \geq 0$. Let u be a nonnegative local weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $p - 1 < s < s_c^* = r(p - 1)/p$, r as in (14). Then for any $B_{R_\infty} \subset B_R \subset B_{R_0} \subseteq \Omega$, the following bound holds*

$$\inf_{x \in B_{R_\infty}} u(x) = \|u\|_{-\infty, R_\infty} \geq \frac{I_{-\infty, q}}{I_{q_0, q}} \frac{\|u\|_{q_0, R}}{|B_R|^{\frac{1}{q_0}}},$$

with $\frac{r[s - (p - 1)]}{r - p} = \bar{q} < q_0 < \frac{r}{p}(p - 1)$

where $q \in (0, \underline{q} \wedge q_0]$ with

$$(54) \quad \underline{q} := \underline{q}_e = \frac{(p - 1)^{\frac{2}{p}} 2^{\frac{(d-1)(p-1)}{p}}}{p \omega_d^2 d^2 e},$$

$I_{-\infty, q}$ as in (46), $q < \underline{q}$ and $\varepsilon = e$ there, and $I_{q_0, q}$ as in (50)-(51).

Proof. It is sufficient to combine the local lower bound proved in Theorem 5.1, with $\varepsilon = e$, the reverse Hölder inequality of Proposition 5.1 and put $\rho_1 = R$ there. \square

6 - Harnack inequalities

In this section we combine the upper bounds of Section 4 with the lower bounds of Section 5 to obtain various forms of Harnack inequalities. The general form, valid in the whole range of exponents, is given in Theorem 6.1. As far as we know, the Harnack inequality that we derive for $s > p - 1$ is not stated explicitly in the literature. Unfortunately, the constant of the general Harnack inequality of Theorem 6.1 depends on u through a quotient of L^q norms. Such quotient simplifies to a constant in some cases and gives clean versions of the Harnack inequality (i.e. the constant does not depend on u); this happens in the subcritical range, i.e. when $0 < s \leq p - 1$, cf. Theorem 6.2, or in the supercritical range $p - 1 < s < s_c^*$, cf. Theorem 6.3. In the range $s_c^* < s < r - 1$, we are not able to prove such clean forms of Harnack inequalities, and we conjecture that the dependence on some L^q norm of the solution can not be avoided, as already mentioned in the introduction. The fact that the “constant” involved has to depend on u when $s_c^* \leq s < r - 1$ is confirmed by the results of [6, 7, 9, 8, 26, 27], [26] applied to separation of variable solutions of parabolic problems, see also [27]. This is also related to the fact that, in the range $s_c^* \leq s < r - 1$, there may exist (very weak, when $p = 2$) singular solutions, cf. [24, 40, 42, 43, 44, 45].

When $s < p^* - 1$, in the case $r = p^*$, so $p < d$, bounded weak solutions are known to be $C^{1,\alpha}$, see [25], and the $C^{1,\alpha}$ modulus of continuity depends on the local L^∞ -norm of the solution or on the constant in the Harnack inequality. Therefore having absolute bounds (independent of u) for the solution or for the Harnack constant, allows to have absolute bounds for the $C^{1,\alpha}$ modulus of continuity. What we show here, is that the $C^{1,\alpha}$ modulus of continuity is independent of the solution when $s < s_c^* < p^* - 1$, while it depends on (some L^q -norms of) the solution when $s_c^* < s < p^* - 1$. If one wants to have a $C^{1,\alpha}$ modulus of continuity independent of u also when $s > s_c^*$, one has to add some extra hypothesis on the solution, and this will be done in the next section, for the special class of stable solutions.

Theorem 6.1 (Harnack inequality for $0 \leq s < s_c$). *Let $\Omega \subset \mathbb{R}^d$. Let u be a nonnegative local weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $p > 1$, $\lambda \geq 0$, $0 \leq s < s_c := r - 1$, r as in (14). Then for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$ and $\varepsilon > 0$, we assume*

$$0 < q_0 \leq \underline{q}_c := \frac{(p-1)^{\frac{2}{p}} 2^{\frac{(d-1)(p-1)}{p}}}{p \omega_d^2 d [e(d-1) + \varepsilon]}, \quad q > \bar{q} := \frac{r[s - (p-1)]_+}{r-p}.$$

Moreover, if $0 < q < s_c^* = r(p-1)/p$ we also assume

$$\frac{\log \left(\frac{r(p-1)(r-p) - rp[s - (p-1)]_+}{qp(r-p) - rp[s - (p-1)]_+} \right)}{\log \frac{r}{p}} \quad \text{is not an integer.}$$

Then the following bound holds true

$$(55) \quad \sup_{x \in B_{R_\infty}} u(x) \leq \mathcal{H}_s[u] \inf_{x \in B_{R_\infty}} u(x)$$

where $\mathcal{H}_s[u]$ depends on u through some local norms as follows

$$\begin{aligned} \mathcal{H}_s[u] &= \mathcal{H}_s[u](d, p, r, R_0, R_\infty, q, q_0, \varepsilon) \\ &= \frac{I_{\infty, q}}{I_{-\infty, q_0}} \frac{\left(\int_{B_{R_0}} u^q \, dx \right)^{\frac{1}{q}}}{\left(\int_{B_{R_0}} u^{q_0} \, dx \right)^{\frac{1}{q_0}}} \frac{\left(\int_{B_{R_0}} u^q \, dx \right)^{\frac{r[s - (p-1)]_+}{q(r-p)q - r[s - (p-1)]_+}}}{\left[\int_{B_{R_\infty}} u^{[s - (p-1)]_+} \, dx \right]^{\frac{r}{(r-p)q - r[s - (p-1)]_+}}} \end{aligned}$$

with $I_{\infty, q}$ as in (26) and $I_{-\infty, q_0}$ as in (46) with $q = q_0$.

Proof. The local upper estimates of Theorems 4.1, give for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$,

$$\begin{aligned}
 \sup_{x \in B_{R_\infty}} u(x) &= \|u\|_{\infty, R_\infty} \\
 (56) \quad &\leq I_{\infty, q} \left[\frac{\left(\int_{B_{R_0}} u^q \, dx \right)^{\frac{[s-(p-1)]_+}{q}}}{\int_{B_{R_\infty}} u^{[s-(p-1)]_+} \, dx} \right]^{\frac{r}{(r-p)q - r[s-(p-1)]_+}} \frac{\|u\|_{q, R_0}}{|B_{R_0}|^{\frac{1}{q}}},
 \end{aligned}$$

for any $q > \bar{q}$, $I_{\infty, q}$ given by (26) and when $0 < q < s_c^*$ we require the additional condition (28). Moreover Theorem 5.1 states that, for any $\varepsilon > 0$,

$$(57) \quad \inf_{x \in B_{R_\infty}} u(x) \frac{|B_{R_0}|^{\frac{1}{q_0}}}{I_{-\infty, q_0} \|u\|_{q_0, R_0}} \geq 1,$$

$I_{-\infty, q_0}$ given by (46) with $q = q_0$ there and

$$0 < q_0 \leq \frac{(p-1)^{\frac{2}{p}} 2^{\frac{(d-1)(p-1)}{p}}}{p \omega_d^2 d [e(d-1) + \varepsilon]} = q_\varepsilon.$$

Combining (56) and (57) we obtain the desired result. □

Theorem 6.2 (Harnack inequality, $0 \leq s \leq p-1$). *Let $\Omega \subset \mathbb{R}^d$ and let $\lambda > 0$. Let u be a nonnegative local weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $0 \leq s \leq p-1$. Then, for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$ the following bound holds true*

$$(58) \quad \sup_{x \in B_{R_\infty}} u(x) \leq \mathcal{H}_s \inf_{x \in B_{R_\infty}} u(x)$$

where

$$\begin{aligned}
 \mathcal{H}_s &= \left\{ \frac{2^d \left[(p-1)^{\frac{2}{p}-1} 2^{\frac{(d-1)(p-1)}{p}} \left(\frac{r}{p}\right)^{n_0 - \frac{1}{p}} + e p \omega_d^2 d \right] \sqrt{\omega_d}}{(p-1)^{\frac{2}{p}-1} 2^{\frac{(d-1)(p-1)}{p}} \left(\frac{r}{p}\right)^{n_0 - \frac{1}{p}} - e(d-1)p \omega_d^2 d} \right\}^{\frac{2}{q_0}} \\
 (59) \quad &\times \left(\frac{r}{p}\right)^{\frac{p^2 r}{q_0(r-p)^2}} \frac{R_0^{\frac{2d}{q_0} + \frac{rd}{q_0(r-p)}} \omega_d^{\frac{1}{q_0}}}{R_\infty^{(d+p)\frac{r}{q_0(r-p)}} (R_0 - R_\infty)^{\frac{2pr}{q_0(r-p)}}} \\
 &\times \left\{ S_p^{2p} [2^{p-1} R_\infty^p + (R_0 - R_\infty)^p] 2^{\frac{pr}{r-p}} q_0^p c_1 \right\}^{\frac{r}{q_0(r-p)}} \\
 &\times \left[\frac{A_{s,0} r}{p q_0} \left(\frac{R_0 - R_\infty}{R_\infty}\right)^p + c_2 \left(\frac{R_0 - R_\infty}{R_\infty}\right)^p + \frac{2^{p-1} c_1^{p-1}}{c_0^p} \right]^{\frac{r}{q_0(r-p)}}
 \end{aligned}$$

with

$$q_0 = \left(\frac{p}{r}\right)^{n_0 - \frac{1}{p}}(p - 1), \quad n_0 = \left\lceil \frac{\log \frac{p \omega_d^2 d e(d-1)}{(p-1)^{\frac{2}{p}-1} 2^{\frac{(d-1)(p-1)}{p}}}}{\log \frac{r}{p}} + \frac{1}{p} \right\rceil + 1,$$

$$\frac{1}{c_0^p} \leq \left(\frac{r}{(r-p)q_0}\right)^p, \quad c_1 = \frac{r^{\frac{1}{p}}}{r^{\frac{1}{p}} - p^{\frac{1}{p}}},$$

$$c_2 = \max \left\{ \frac{|pq_0 - r(p-1)| r^p}{(pq_0)^{p+1}}, \left(\frac{p}{p-1}\right)^p \frac{1}{(p+1)^{p+1}} \right\}$$

and

$$A_{s,0} = \frac{\lambda}{p^{p-1}} R_0^p \quad \text{if } s = p - 1 \quad \text{and} \quad A_{s,0} = 2^{p-1+d} \quad \text{if } s \neq p - 1.$$

Proof. The goal of the proof is to simplify the quotient of L^q -norms in the expression of the constant $\mathcal{H}_s[u]$ of the Harnack inequality (55). Since we are dealing with the range $0 \leq s \leq p - 1$, we can choose any $q > 0$, hence we can let

$$0 < q = \underline{q}_\varepsilon = q_0 = q_0(\varepsilon) = \frac{(p-1)^{\frac{2}{p}} 2^{\frac{(d-1)(p-1)}{p}}}{p \omega_d^2 d [e(d-1) + \varepsilon]}$$

with i.p. $\left\lceil \frac{\log \frac{r(p-1)}{qp}}{\log \frac{r}{p}} \right\rceil \neq \frac{\log \frac{r(p-1)}{qp}}{\log \frac{r}{p}}.$

In fact, we shall arrive, with a suitable choice of the parameter ε , at a value of q_0 smaller than $r(p-1)/p$, so that the request $\log \frac{r(p-1)}{qp} / \log \frac{r}{p}$ not be integer is necessary. The last condition means $q_0(\varepsilon) \neq (p/r)^{n-1}(p-1)$ for all $n \in \mathbb{N}$ and this is possible since we can always choose ε

$$0 < \varepsilon = \bar{\varepsilon} := \frac{(p-1)^{\frac{2}{p}-1} 2^{\frac{(d-1)(p-1)}{p}}}{p \omega_d^2 d} \left(\frac{r}{p}\right)^{n_0 - \frac{1}{p}} - e(d-1)$$

so that $q_0 = \left(\frac{p}{r}\right)^{n_0 - \frac{1}{p}}(p - 1),$

where n_0 is the first integer n such that $\varepsilon(n) > 0$, which is

$$n_0 = \left\lceil \frac{\log \frac{p \omega_d^2 d e(d-1)}{(p-1)^{\frac{2}{p}-1} 2^{\frac{(d-1)(p-1)}{p}}}}{\log \frac{r}{p}} + \frac{1}{p} \right\rceil + 1.$$

The constants become in this case

$$I_{\infty,q} = I_{\infty,q_0} = \left\{ \frac{|B_{R_0}|^{\frac{r-p}{r}+1}}{|B_{R_\infty}|} \frac{S_p^p q_0^p p^p c_1}{r^p (R_0 - R_\infty)^p} \left[\frac{A_{s,0} r}{p q_0} \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p \right. \right. \\ \left. \left. + \left(\frac{R_0 - R_\infty}{R_\infty} \right)^p c_2 + \frac{c_1^{p-1} 2^{p-1}}{c_0^p} \right] \left(\frac{r}{p} \right)^{\frac{rp}{r-p}} \right\}^{\frac{r}{q_0(r-p)}}$$

where $A_{s,0}$ is given by (27),

$$\frac{1}{c_0^p} \leq \left(\frac{r}{q_0(r-p)} \right)^p, \\ c_2 = \max \left\{ \frac{|pq_0 - r(p-1)| r^p}{(p q_0)^{p+1}}, \left(\frac{p}{p-1} \right)^p \frac{1}{(p+1)^{p+1}} \right\}$$

and since $q_0 < r(p-1)/p$

$$c_1 = \max_{i=0,1} \frac{\left(\frac{r}{p} \right)^{j_0+i-1} q_0}{\left| \left(\frac{r}{p} \right)^{j_0+i-1} q_0 - (p-1) \right|} = \max_{i=0,1} \frac{\left(\frac{r}{p} \right)^{i+\frac{1}{p}}}{\left(\frac{r}{p} \right)^{i+\frac{1}{p}} - 1} = \frac{r^{\frac{1}{p}}}{r^{\frac{1}{p}} - p^{\frac{1}{p}}}$$

since

$$j_0 = \text{i.p.} \left[\frac{\log \frac{r(p-1)}{q_0 p}}{\log \frac{r}{p}} \right] = \text{i.p.} \left[1 + \frac{\log \frac{p-1}{q_0}}{\log \frac{r}{p}} \right] = \text{i.p.} \left[n_0 + 1 - \frac{1}{p} \right] = n_0 + 1.$$

Moreover

$$I_{-\infty,q_0} = \left[\frac{(R_0 - R_\infty) R_\infty}{R_0^{\frac{d(r-p)}{rp}}} \right]^{\frac{pr}{q_0(r-p)}} \\ \times \left\{ S_p^p [2^{p-1} R_\infty^p + (R_0 - R_\infty)^p] 2^{\frac{pr}{r-p}} \right\}^{-\frac{r}{q_0(r-p)}} \\ \times \left\{ \frac{(p-1)^{\frac{2}{p}-1} 2^{\frac{(d-1)(p-1)}{p}} \left(\frac{r}{p} \right)^{n_0 - \frac{1}{p}} - e(d-1)p\omega_d^2 d}{2^d \left[(p-1)^{\frac{2}{p}-1} 2^{\frac{(d-1)(p-1)}{p}} \left(\frac{r}{p} \right)^{n_0 - \frac{1}{p}} + e p \omega_d^2 d \right] \sqrt{\omega_d}} \right\}^{\frac{2}{q_0}}.$$

Hence we get the expression of $\mathcal{H}_s = I_{\infty,q}/I_{-\infty,q_0}$ as in (59). □

Unfortunately, when $s > p - 1$ we can not join the upper and the lower bound so easily, we need a further iteration.

Theorem 6.3 (Harnack Inequalities when $p - 1 < s < s_c^*$). *Let $\Omega \subset \mathbb{R}^d$. Let u be a nonnegative local weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $p > 1$, $\lambda \geq 0$ and $p - 1 < s < s_c^* = r(p - 1)/p$. Then for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$ there exists an explicit constant $\mathcal{H}_s > 0$ such that*

$$(60) \quad \sup_{x \in B_{R_\infty}} u(x) \leq \mathcal{H}_s \inf_{x \in B_{R_\infty}} u(x)$$

where \mathcal{H}_s does not depend on u , and is given by

$$(61) \quad \mathcal{H}_s = I_{\infty, q_0} \left(\frac{I_{q_0, q}}{I_{-\infty, q}} \right)^{\frac{q_0(r-p)}{q_0(r-p) - r[s-(p-1)]}}$$

with $q \in (0, \underline{q} \wedge q_0]$, \underline{q} and $I_{-\infty, q}$ are given in (54) and (46) respectively, $I_{q_0, q}$ in (50) and (51) and I_{∞, q_0} in (26) with $q = q_0$; moreover, since $q < s_c^*$ we require the additional condition (28).

Proof. Let $B_{R_\infty} \subset B_R \subset B_{R_0} \subseteq \Omega$, then by Theorem 5.2 we have

$$(62) \quad \frac{\|u\|_{q_0, R}}{|B_R|^{\frac{1}{q_0}}} \leq \frac{I_{q_0, q}}{I_{-\infty, q}} \inf_{x \in B_{R_\infty}} u(x),$$

with $\bar{q} < q_0 < r(p - 1)/p$, $q \in (0, \underline{q} \wedge q_0]$, \underline{q} as in (54), $I_{q_0, q}$ as in (50) and (51) and $I_{-\infty, q}$ as in (46). Moreover Theorem 4.1, applied with $R_0 = R$ and $q = q_0 > \bar{q}$, gives

$$\begin{aligned} \sup_{x \in B_{R_\infty}} u(x) &= \|u\|_{\infty, R_\infty} \leq I_{\infty, q_0} \frac{\|u\|_{q_0, R}}{|B_R|^{\frac{1}{q_0}}} \left(\frac{\|u\|_{q_0, R}^{[s-(p-1)]}}{|B_R|^{\frac{[s-(p-1)]}{q_0}}} \right)^{\frac{r}{(r-p)q_0 - r[s-(p-1)]}} \\ &\quad \times \left(\frac{|B_{R_\infty}|}{\int_{B_{R_\infty}} u^{[s-(p-1)]} dx} \right)^{\frac{r}{(r-p)q_0 - r[s-(p-1)]}} \\ &\leq I_{\infty, q_0} \frac{\|u\|_{q_0, R}}{|B_R|^{\frac{1}{q_0}}} \\ &\quad \times \left(\frac{\|u\|_{q_0, R}}{|B_R|^{\frac{1}{q_0}}} \frac{1}{\inf_{x \in B_{R_\infty}} u(x)} \right)^{\frac{r[s-(p-1)]}{(r-p)q_0 - r[s-(p-1)]}}, \end{aligned}$$

for any $q_0 > r[s - (p - 1)]/(r - p)$ and I_{∞, q_0} as in (26). Therefore, using twice the lower bound (62) in the previous inequality, we conclude the proof of the theorem. □

7 - Local absolute bounds

The interest of having absolute upper bounds for solutions of nonlinear elliptic equations is related to several aspects of the theory of such equations. If we have at our disposal local absolute upper bounds, then the constant in the general Harnack inequality of Theorem 6.1 can be independent of u and also the $C^{1,\alpha}$ modulus of continuity will be independent of u , as already discussed at the beginning of Section 6. The absolute estimates that we present here have a local nature, which means that they are independent of the boundary conditions, which can be of Dirichlet, Neumann, Robin type, or also “large”, i.e. $u = +\infty$ on the boundary. Such absolute bounds have many more applications, for example, they may imply Liouville-type Theorems on \mathbb{R}^d [32, 52], they imply existence of large solutions, and the fact that the constant is explicit is really useful although not always indispensable. For the homogeneous Dirichlet problem for semilinear equations (namely for $p = 2$), absolute upper bounds, sometimes called universal bounds, have been proved by many authors, [11, 23, 31, 32, 46, 52], but in those papers the constant was not quantitative, and to our knowledge it can not be made quantitative with the proofs presented there. An effort to provide quantitative global absolute bounds for this Dirichlet problem has been done in [5].

In this section we first prove absolute upper and lower bounds for weak solutions, in the range $p - 1 < s < s_c^*$ and $0 < s \leq p - 1$ respectively. Next we want to obtain quantitative absolute upper bounds for $s > s_c^*$, which are known to be false in the whole class of weak (or very weak when $p = 2$) solutions, in view of the existence of singular solutions, as already mentioned in the Introduction, cf. [24, 40, 42, 43, 44, 45]; therefore we have to pass to a special class of solutions, the so-called stable solutions [13, 14, 16, 22, 28, 29], for which we can bound absolutely from above the $L^{\bar{m}}$ -norm of the solution, for \bar{m} sufficiently large, and we combine such bounds with the upper bounds of type II of Theorem 4.4 to get our quantitative absolute upper bounds for stable solutions. We can cover the whole range of $s \geq 0$ only for small spatial dimensions, namely $d \leq \frac{p(p+3)}{p-1}$; for larger dimensions, it appears a new exponent $r - 1 < s_{JL} < \infty$, the so-called Joseph-Lundgren exponent, and the absolute bounds hold only until that exponent.

7.1 - Local absolute bounds for $s < s_c^*$

In this section we will prove local absolute lower bounds when $0 \leq s < p - 1$ and a local absolute upper bounds when $p - 1 < s < s_c^*$ as a consequence of the Harnack inequalities of the previous section together with the Caccioppoli estimate (12).

Theorem 7.1 (Local absolute bounds). *Let $\Omega \subset \mathbb{R}^d$. Let u be a nonnegative local weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $p > 1$, $\lambda \geq 0$ and $0 \leq s < s_c^* = r(p-1)/p$, r as in (14). Then for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$ there exists a constant \mathcal{H}_s that does not depend on u , such that*

$$\sup_{x \in B_{R_\infty}} u(x) \leq \mathcal{H}_s \left(\frac{p^{p-1} 2^{p-1} R_0^d}{\lambda (R_0 - R_\infty)^p R_\infty^d} \right)^{\frac{1}{s-(p-1)}} \quad \text{if } p-1 < s < s_c^*$$

with \mathcal{H}_s given by (61), and, if $u \neq 0$ on B_{R_0}

$$\inf_{x \in B_{R_\infty}} u(x) \geq \mathcal{H}_s^{-1} \left(\frac{\lambda (R_0 - R_\infty)^p R_\infty^d}{p^{p-1} 2^{p-1} R_0^d} \right)^{\frac{1}{(p-1)-s}} \quad \text{if } 0 \leq s < p-1,$$

with \mathcal{H}_s given by (59).

Proof. First, we note that the Caccioppoli estimate (12), with $\rho_1 = R_\infty$ and $\rho_0 = R_0$, implies when $s > p-1$

$$\begin{aligned} \inf_{x \in B_{R_\infty}} u(x) &\leq \left(\frac{1}{|B_{R_\infty}|} \int_{B_{R_\infty}} u^{s-(p-1)} \, dx \right)^{\frac{1}{s-(p-1)}} \\ &\leq \left(\frac{p^{p-1} 2^{p-1} R_0^d}{\lambda (R_0 - R_\infty)^p R_\infty^d} \right)^{\frac{1}{s-(p-1)}}. \end{aligned}$$

Moreover, since we are assuming $u \neq 0$ on B_{R_0} , if $0 \leq s < p-1$, (12), applied always with $\rho_1 = R_\infty$ and $\rho_0 = R_0$, gives

$$\left(\frac{\lambda (R_0 - R_\infty)^p R_\infty^d}{p^{p-1} 2^{p-1} R_0^d} \right)^{\frac{1}{(p-1)-s}} \leq \left(\frac{|B_{R_\infty}|}{\int_{B_{R_\infty}} u^{s-(p-1)} \, dx} \right)^{\frac{1}{(p-1)-s}} \leq \sup_{x \in B_{R_\infty}} u(x).$$

The above estimates can be now combined with the corresponding Harnack inequalities (60) and (58) to obtain the desired bounds in both cases. □

7.2 - Local absolute bounds for stable solutions. The supercritical case

In this section we establish local upper bounds for stable solutions. From now on, we assume $p \geq 2$. The results can be proved also in the case $1 < p < 2$, but we need some modifications in the definitions of stable solutions and in the proofs. We have decided to deal with $p \geq 2$ in order to simplify the exposition. When $1 < p < 2$ we refer to [14] and references therein. Let us mention that the proof that we give here is a modification of an idea originally due to A. Farina, see [16, 28, 29]; see also [13, 14]

for an alternative approach. Our proof is slightly different from [16, 28, 29] and provides explicit constants.

Definition 7.1. A function u is a *local stable solution* to $-\Delta_p u = \lambda u^s$, if and only if $0 \leq u \in W_{loc}^{1,p}(\Omega)$ and satisfies

$$(63) \quad \int_{\Omega} \left\{ |\nabla u|^{p-2} \left[|\nabla \varphi|^2 + (p-2) \left(\nabla \varphi \cdot \frac{\nabla u}{|\nabla u|} \right)^2 \right] - \lambda s u^{s-1} \varphi^2 \right\} dx \geq 0$$

for all bounded $\varphi \in C_c^1(\Omega)$.

We recall that the stability condition translates into the fact that the second variation of the energy functional is nonnegative, see [13, 14, 16, 22, 28, 29] for a more detailed study of stable solutions related to this kind of problem.

Remark 7.1. From the stability condition (63) we immediately obtain

$$(64) \quad \lambda s \int_{\Omega} u^{s-1} \varphi^2 dx \leq (p-1) \int_{\Omega} |\nabla u|^{p-2} |\nabla \varphi|^2 dx,$$

for all bounded $\varphi \in C_c^1(\Omega)$.

Now we have the following estimate for nonnegative stable solutions, which we already know to be bounded, by the upper estimates of the previous sections.

Lemma 7.1. *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain and let u be a local non-negative bounded stable weak solution to $-\Delta_p u = \lambda u^s$ in Ω , $\lambda \geq 0$ and $s > p - 1$. Then the following estimate holds true for any $\alpha > 0$, $\delta, \varepsilon \in (0, 1]$ and any test function $\phi \in C_c^1(\Omega)$, $\phi > 0$*

$$(65) \quad \begin{aligned} \lambda s \int_{\Omega} u^{s+\alpha} \phi^{\frac{s+\alpha}{s-(p-1)}} dx &\leq \left[(p-1)(1+\varepsilon) \left(\frac{\alpha+1}{2} \right)^2 \right. \\ &\quad \left. + \frac{p-1}{4} \left(1 + \frac{1}{\varepsilon} \right) \frac{(s+\alpha)^2}{[s-(p-1)]^2} \frac{\delta(p-2)}{p} \right] \\ &\quad \times \frac{p^p}{[\alpha+(p-1)]^p} \int_{\Omega} \left| \nabla \left(u^{\frac{\alpha+(p-1)}{p}} \right) \right|^p \phi^{\frac{s+\alpha}{s-(p-1)}} dx \\ &\quad + \frac{p-1}{4} \left(1 + \frac{1}{\varepsilon} \right) \frac{(s+\alpha)^2}{[s-(p-1)]^2} \frac{2}{p \delta^{\frac{p-2}{p}}} \\ &\quad \times \int_{\Omega} u^{\alpha+(p-1)} |\nabla \phi|^p \phi^{\frac{s+\alpha}{s-(p-1)-p}} dx. \end{aligned}$$

Proof. Let $0 < \phi \in C_c^1(\Omega)$. Using a standard approximation technique (recall that u is bounded), we can use as test function $\varphi^2 := u^{\alpha+1}\phi^\gamma$, with

$$\gamma := \frac{s + \alpha}{s - (p - 1)} > 1, \quad \text{since } \alpha > 0$$

in (64) and Young inequality with $\varepsilon > 0$, we get

$$\begin{aligned} \lambda s \int_{\Omega} u^{s+\alpha} \phi^\gamma \, dx &\leq (p-1)(1+\varepsilon) \left(\frac{\alpha+1}{2}\right)^2 \int_{\Omega} u^{\alpha-1} |\nabla u|^p \phi^\gamma \, dx \\ &\quad + \frac{p-1}{4} \left(1 + \frac{1}{\varepsilon}\right) \gamma^2 \int_{\Omega} |\nabla u|^{p-2} u^{\alpha+1} |\nabla \phi|^2 \phi^{\gamma-2} \, dx. \end{aligned}$$

Again using Young inequality with $\delta > 0$ and exponents $p/2, p/(p-2)$ we obtain, for the second terms in the right hand side of the previous inequality, the following estimate

$$\begin{aligned} \int_{\Omega} |\nabla u|^{p-2} u^{\alpha+1} |\nabla \phi|^2 \phi^{\gamma-2} \, dx &\leq \frac{\delta(p-2)}{p} \int_{\Omega} |\nabla u|^p u^{\alpha-1} \phi^\gamma \, dx \\ &\quad + \frac{2}{p \delta^{\frac{p-2}{2}}} \int_{\Omega} u^{\alpha+(p-1)} |\nabla \phi|^p \phi^{\gamma-p} \, dx. \end{aligned}$$

Combining the previous estimates and noticing that

$$\int_{\Omega} |\nabla u|^p u^{\alpha-1} \phi^\gamma \, dx = \frac{p^p}{[\alpha + (p-1)]^p} \int_{\Omega} \left| \nabla \left(u^{\frac{\alpha+(p-1)}{p}} \right) \right|^p \phi^\gamma \, dx$$

we arrive at the desired result. \square

Combining the previous estimate, coming from the stability condition (64), and the following form of the energy estimate (5),

$$\begin{aligned} &\frac{p^p}{[\alpha + (p-1)]^p} \int_{\Omega} \left| \nabla \left(u^{\frac{\alpha+(p-1)}{p}} \right) \right|^p \phi^{\frac{s+\alpha}{s-(p-1)}} \, dx \\ &\leq \frac{\lambda p [s - (p-1)]}{p \alpha [s - (p-1)] - (s + \alpha) \bar{\varepsilon} (p-1)} \int_{\Omega} u^{\alpha+s} \phi^{\frac{s+\alpha}{s-(p-1)}} \, dx \\ (66) \quad &\quad + \frac{s + \alpha}{\bar{\varepsilon}^{p-1} \{p \alpha [s - (p-1)] - (s + \alpha) \bar{\varepsilon} (p-1)\}} \\ &\quad \times \int_{\Omega} u^{\alpha+(p-1)} |\nabla \phi|^p \phi^{\frac{s+\alpha}{s-(p-1)-p}} \, dx \end{aligned}$$

for any

$$0 < \bar{\varepsilon} < \frac{p\alpha[s - (p - 1)]}{(s + \alpha)(p - 1)}, \quad \alpha > 0 \quad \text{and} \quad s > p - 1.$$

For the proof of the above inequality we have to follow the proof of Lemma 2.1 and change slightly the test function (we have to use $u^\alpha \phi^{\frac{s+\alpha}{s-(p-1)}}$ instead of $(u + \delta)^\alpha \phi$). We get the following.

Lemma 7.2. *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain and let u be a local non-negative stable weak solution to $-\Delta_p u = \lambda u^s$ in Ω , $\lambda > 0$ and $s > p - 1$. Then the following estimate holds true for any*

$$(67) \quad 0 < \alpha < \bar{\alpha} := \frac{2s - (p - 1) + 2\sqrt{s^2 - s(p - 1)}}{p - 1},$$

and any test function $\phi \in C_c^1(\Omega)$, $\phi > 0$

$$(68) \quad \int_{\Omega} u^{s+\alpha} \phi^{\frac{s+\alpha}{s-(p-1)}} \, dx \leq c_4 \int_{\Omega} u^{\alpha+(p-1)} |\nabla \phi|^p \phi^{\frac{s+\alpha}{s-(p-1)-p}} \, dx,$$

where c_4 is a positive constant that depends on s, p, λ and α .

Proof. Using (66) to estimate the second term in the right hand side of (65) and operating some simple manipulations, we get for any $\alpha > 0$

$$k_1 \int_{\Omega} u^{\alpha+s} \phi^{\frac{s+\alpha}{s-(p-1)}} \, dx \leq k_2 \int_{\Omega} u^{\alpha+(p-1)} |\nabla \phi|^p \phi^{\frac{s+\alpha}{s-(p-1)-p}} \, dx,$$

where $k_i = k_i(\alpha, \delta, \varepsilon, \bar{\varepsilon}, p, s) > 0$, $i = 1, 2$, for any $\varepsilon > 0$, $\delta > 0$ and $0 < \bar{\varepsilon} < \frac{p\alpha[s - (p - 1)]}{(s + \alpha)(p - 1)}$, more precisely,

$$k_1 = \frac{\lambda p[s - (p - 1)]}{p[s - (p - 1)]\alpha - (s + \alpha)\bar{\varepsilon}(p - 1)} \left\{ s \left(\alpha - \bar{\varepsilon} \frac{(p - 1)(\alpha + s)}{p[s - (p - 1)]} \right) - \left[(p - 1)(1 + \varepsilon) \left(\frac{\alpha + 1}{2} \right)^2 + \delta \left(1 + \frac{1}{\varepsilon} \right) \frac{(s + \alpha)^2}{[s - (p - 1)]^2} \frac{(p - 2)(p - 1)}{4p} \right] \right\}.$$

First, we notice that letting $\varepsilon = \bar{\varepsilon} = \delta = 0$ in the above expression, gives a positive number whenever $0 < \alpha < \bar{\alpha}$. As a consequence of the explicit continuous dependence on the parameters ε, δ and $\bar{\varepsilon}$, we can always fix them sufficiently small so that $k_1 > 0$ when $0 < \alpha < \bar{\alpha}$, therefore the constant $c_4 = k_2/k_1 > 0$. \square

Theorem 7.2. *Let $\Omega \subset \mathbb{R}^d$. Let u be a local nonnegative stable weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $\lambda \geq 0$ and $s > p - 1$. Then for any $B_{\rho_1} \subset B_{\rho_0} \subset\subset \Omega$ there exists a constant that does not depend on u such that*

$$(69) \quad \|u\|_{s+\alpha, \rho_1} \leq c_5$$

for any

$$(70) \quad 0 < \alpha < \bar{\alpha} := \frac{2s - (p - 1) + 2\sqrt{s^2 - s(p - 1)}}{p - 1},$$

and

$$(71) \quad c_5 = c_4^{\frac{1}{s-(p-1)}} \left[\frac{2^{p-1} p^p}{(\rho_0 - \rho_1)^p} \right]^{\frac{1}{s-(p-1)}} \rho_0^{\frac{d}{s+\alpha}} \omega_d^{\frac{1}{s+\alpha}}.$$

Proof. The result follows from the previous lemma, Hölder inequality and using the test function defined in Lemma 2.2. Indeed, by Hölder inequality with exponents $(s + \alpha)/[\alpha + (p - 1)]$ and $(s + \alpha)/[s - (p - 1)]$, applied to the right hand side of (68), we have

$$\begin{aligned} \int_{\Omega} u^{s+\alpha} \phi^{\frac{s+\alpha}{s-(p-1)}} dx &\leq c_4 \left[\int_{\Omega} u^{s+\alpha} \phi^{\frac{s+\alpha}{s-(p-1)}} dx \right]^{\frac{\alpha+(p-1)}{\alpha+s}} \\ &\quad \times \left[\int_{\Omega} \left(\frac{|\nabla \phi|^p}{\phi^{p-1}} \right)^{\frac{s+\alpha}{s-(p-1)}} dx \right]^{\frac{s-(p-1)}{s+\alpha}}. \end{aligned}$$

Hence, we arrive to the desired results simplifying and choosing ϕ as in Lemma 2.2. □

The Joseph-Lundgren exponent s_{JL} . The above theorem proves absolute bounds for some local $L^{\bar{m}}$ -norm, and we would like to have \bar{m} sufficiently large, namely

$$\bar{m} > \frac{r[s - (p - 1)]}{r - p} = \bar{q}$$

to be able to combine the above absolute bounds (69) with the upper bounds of type II of Theorem 4.4. Letting then $\bar{m} = s + \alpha$, with α satisfying the condition (70), we have that

$$\bar{q} < \bar{m} = s + \alpha < s + \bar{\alpha} = s + \frac{2s - (p - 1) + 2\sqrt{s^2 - s(p - 1)}}{p - 1}$$

where we take $r = p^* = pd/(d - p)$, the Sobolev exponent, i.e. we are in the case

$p < d$. Notice that when $p \geq d$, we can take r large enough and the above condition is always satisfied. In the case under consideration, namely $1 < p < d$, the above condition is satisfied by all the s in some interval, more precisely, there exists an exponent s_{JL} such that for all $s \in [0, s_{JL})$ we have

$$\|u\|_{\overline{m}, R_1} \leq c_5, \quad \text{with } \overline{m} > \frac{r[s - (p - 1)]}{r - p}$$

where c_5 is given in (71). Moreover, we call the exponent s_{JL} the Joseph-Lundgren exponent and it has the explicit form

$$(72) \quad s_{JL} := \frac{[(p - 1)d - p]^2 + p^2(p - 2) - p^2(p - 1)d + 2p^2\sqrt{(p - 1)(d - 1)}}{(d - p)[(p - 1)d - p(p + 3)]}$$

if $d > [p(p + 3)]/(p - 1)$ and

$$s_{JL} := +\infty \quad \text{if } d \leq \frac{p(p + 3)}{p - 1}.$$

See [29, 30, 35] for more details on the derivation of the Joseph-Lundgren exponent.

All the above discussion can be summarized in the following:

Theorem 7.3 (Local absolute bounds for stable solutions). *Let $\Omega \subset \mathbb{R}^d$. Let u be a local nonnegative stable weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $\lambda \geq 0$ and $p - 1 < s < s_{JL}$. Then for any $B_{R_\infty} \subset B_{R_0} \subseteq \Omega$ there exists a constant that does not depend on u such that, for any $\overline{m} > r[s - (p - 1)]/(r - p)$,*

$$(73) \quad \|u\|_{\infty, R_\infty} \leq c_6$$

where

$$(74) \quad c_6 = \frac{A_{\overline{m}}^{(1)} c_5}{(R_0 - R_\infty)^{\frac{pr}{\overline{m}(r-p)}}} \left[A_{\overline{m}}^{(2)} + A_{\overline{m}}^{(3)} \lambda^{\frac{\overline{m}r}{\overline{m}(r-p) - r[s - (p-1)]}} c_5^{\frac{\overline{m}r[s - (p-1)]}{\overline{m}(r-p) - r[s - (p-1)]}} \right]^{\frac{r}{\overline{m}(r-p)}}$$

and $A_{\overline{m}}^{(i)}$, $i = 1, 2, 3$, are as in Theorem 4.3 with $q = \overline{m}$ and c_5 is given in (71).

Proof. Combine the upper bounds, established in Theorem 4.4, choosing $q = \overline{m}$ there, with the absolute upper bounds (69), with $\rho_0 = R_0$ and $\rho_1 = R_\infty$, to get

$$\|u\|_{\infty, R_\infty} \leq \frac{A_{\overline{m}}^{(1)}}{(R_0 - R_\infty)^{\frac{pr}{\overline{m}(r-p)}}} \left[A_{\overline{m}}^{(2)} + A_{\overline{m}}^{(3)} \lambda^{\frac{\overline{m}r}{\overline{m}(r-p) - r[s - (p-1)]}} c_5^{\frac{\overline{m}r[s - (p-1)]}{\overline{m}(r-p) - r[s - (p-1)]}} \right]^{\frac{r}{\overline{m}(r-p)}} c_5$$

where $A_{\overline{m}}^{(i)}$, $i = 1, 2, 3$, are as in Theorem 4.3 with $q = \overline{m} > \overline{q}$ and c_5 is given in (71). □

8 - Appendix

8.1 - The John-Nirenberg Lemma and reverse Hölder inequalities

First of all we recall a quantitative version of Lemma 7.20 of [33], proved in [4] (see Lemma 4.2 there).

From now on we denote, as usual, by $M^m(\Omega)$ the Marcinkiewicz spaces for any $m > 1$ and by $\mathcal{V}_\mu[g]$ the Riesz potential of a function g , that is

$$\mathcal{V}_\mu[g](x) = \int_{\Omega} \frac{g(y)}{|x - y|^{d(1-\mu)}} dy, \quad \mu \in (0, 1].$$

Lemma 8.1 (A “potential” version of the Moser-Trudinger imbedding). *Let $g \in M^\sigma(\Omega)$ with $\sigma > 1$ and let us suppose $\|g\|_{M^\sigma(\Omega)} \leq K$. Then there exist two constants k_2 and k_3 such that*

$$\int_{\Omega} \exp \left[\frac{|\mathcal{V}_{\frac{1}{\sigma}}[g](x)|}{k_2 K} \right] dx \leq k_3.$$

One can take

$$k_2 > (\sigma - 1)e \quad \text{and} \quad k_3 = |\Omega| + \frac{\text{diam}(\Omega)^d}{\sqrt{2\pi}} \frac{se\omega_d}{k_2 - (\sigma - 1)e}.$$

Now we state a quantitative version of Jonh-Nirenberg lemma for convex domains; for the proof see Lemma 4.3 in [4].

Lemma 8.2 (Jonh-Nirenberg). *Let $g \in W^{1,1}(\Omega)$ where Ω is convex, and suppose there exists a constant K such that*

$$\int_{B_R \cap \Omega} |\nabla g| dx \leq K R^{d-1}, \quad \text{for all balls } B_R \subseteq \Omega.$$

Then the following inequality holds

$$\int_{\Omega} \exp \left[\frac{|g - g_{\Omega}|}{k_0 K} \right] dx \leq k_1,$$

where for any $k_2 > (d - 1)e$

$$k_0 = \frac{d|\Omega|}{\text{diam}(\Omega)^d} k_2, \quad k_1 = \frac{\omega_d \text{diam}(\Omega)^d (k_2 + e)}{k_2 - (d - 1)e} \quad \text{and} \quad g_{\Omega} = \frac{1}{|\Omega|} \int_{\Omega} g dx.$$

The John-Nirenberg Lemma has an important consequence when applied to the function $g = \log(u + \delta)$, $\delta > 0$.

Proposition 8.1 (Reverse Hölder inequalities). *Let $\delta \geq 0$ and let u be a positive measurable function such that $\log(u + \delta) \in W^{1,1}(\Omega)$, where Ω is convex, and suppose there exists a constant K (independent of δ) such that*

$$\int_{B_R \cap \Omega} |\nabla \log(u + \delta)| \, dx \leq K R^{d-1}, \quad \text{for all balls } B_R \subseteq \Omega.$$

Then the following inequality

$$\frac{\|u + \delta\|_{q,\Omega}}{\|u + \delta\|_{-q,\Omega}} \leq k_1^{2/q}$$

holds for any

$$0 < q \leq \frac{1}{k_0 K},$$

where the constants k_i are given in Lemma 8.2.

Proof. See Proposition 4.4 in [4]. □

We conclude this section by showing that reverse Hölder inequalities hold for local solutions to our problem, as a consequence of Caccioppoli estimates (see Corollary 2.1).

Proposition 8.2 (Reverse Hölder inequalities). *Let $\Omega \subset \mathbb{R}^d$ and let $\lambda \geq 0$. Let u be a local weak solution to $-\Delta_p u = \lambda u^s$ in Ω , with $0 \leq s < s_c = r - 1$, r as in (14). Then for any $\varepsilon > 0$, the following inequality holds true for any $\delta \geq 0$ and for any $B_{R_0} \subseteq \Omega$*

$$(75) \quad \left[\frac{\varepsilon}{2^d (e d + \varepsilon)} \right]^{2/q} \frac{\|u + \delta\|_{q,R_0}}{|B_{R_0}|^{\frac{1}{q}}} \leq \frac{\|u + \delta\|_{-q,R_0}}{|B_{R_0}|^{\frac{1}{q}}},$$

$$\text{for all } 0 < q \leq \frac{(p-1)^{\frac{2}{p}} 2^{\frac{(d-1)(p-1)}{p}}}{p \omega_d^2 d [e(d-1) + \varepsilon]} =: \underline{q}_\varepsilon.$$

Proof. The Caccioppoli estimate (11), with $\rho_1 = R$ and $\rho_0 = 2R$, R any positive real number less or equal than R_0 , implies that

$$\int_{B_R} |\nabla \log(u + \delta)|^p \, dx \leq \frac{2^{d+p-1} p^p R^{d-p} \omega_d}{(p-1)^2},$$

hence the hypothesis of the previous proposition are satisfied, more precisely

$$\int_{B_{R_0} \cap B_R} |\nabla \log(u + \delta)| \, dx \leq |B_R|^{1-\frac{1}{p}} \left[\int_{B_R} |\nabla \log(u + \delta)|^p \, dx \right]^{\frac{1}{p}} \\ \leq R^{d-1} \omega_d \frac{2^{\frac{p+d-1}{p}} p}{(p-1)^{\frac{p}{2}}}.$$

Therefore putting $K = \omega_d \frac{2^{\frac{p+d-1}{p}} p}{(p-1)^{\frac{p}{2}}}$, taking an $\varepsilon > 0$ and choosing $k_2 = e(d-1) + \varepsilon$, by the Proposition 8.1, we get the desired result. \square

8.2 - Technical tools

In this section we recall, in order to be complete, some tools that we use in this paper. The first lemma concerns the geometric convergence of some sequences of real numbers.

Lemma 8.3 (Numerical Iteration). *Let $Y_n \geq 0$ be a sequence of numbers such that*

$$Y_n \leq I_{n-1}^{\sigma \theta^{n-1}} Y_{n-1} \quad \text{with} \quad I_{n-1} \leq I_0 C^{n-1}$$

for some $\sigma, I_0, C > 0, \theta \in (0, 1)$. Then $\{Y_n\}$ is a bounded sequence and one has

$$Y_\infty := \limsup_{n \rightarrow +\infty} Y_n \leq I_0^{\frac{\sigma}{1-\theta}} C^{\frac{\sigma \theta}{1-\theta^2}} Y_0.$$

Proof. See for example Lemma 7.1 of [34]. \square

The following lemma is due to E. De Giorgi and its proof is contained in several books and papers, see for example [34], Lemma 6.1.

Lemma 8.4 (De Giorgi). *Let Z be a bounded nonnegative function in the interval $[t_0, t_1]$. Assume that for $t_0 \leq t < s \leq t_1$ we have*

$$Z(t) \leq \theta Z(s) + \frac{A}{(s-t)^\alpha},$$

with $A \geq 0, \alpha > 0$ and $0 \leq \theta < 1$. Then

$$Z(t_0) \leq \frac{A c(\alpha, \lambda, \theta)}{(t_1 - t_0)^\alpha}$$

where

$$c(\alpha, \lambda, \theta) = \frac{1}{(1 - \lambda)^\alpha \left(1 - \frac{\theta}{\lambda^\alpha}\right)} \text{ for any } \lambda \in (\theta^{\frac{1}{\alpha}}, 1).$$

This lemma has an important consequence, indeed it is necessary to obtain extending local upper bounds (see Section 4). More precisely, it allows to prove that, if a reverse Hölder inequality holds for some $0 < q_1 < q_2 \leq \infty$, then it holds for any $0 < q < q_2 \leq \infty$.

Lemma 8.5. *Assume that the following bound holds true for some $0 < q_1 < q_2 < \infty$ and for any $R_\infty \leq \rho_1 < \rho_0 \leq R_0$,*

$$\|u\|_{q_2, \rho_1} \leq \frac{K}{(\rho_0 - \rho_1)^\gamma} \|u\|_{q_1, \rho_0}.$$

Then we have that for all $0 < q \leq q_1 < q_2 < \infty$

$$\|u\|_{q_2, R_\infty} \leq 3 \cdot 2^{\frac{q_2(q_1 - q)}{q(q_2 - q_1)}} \left[\left(4^\gamma \frac{q_1(q_2 - q)}{q(q_2 - q_1)} \right)^\gamma \frac{K}{(R_0 - R_\infty)^\gamma} \right]^{\frac{q_1(q_2 - q)}{q(q_2 - q_1)}} \|u\|_{q, R_0}.$$

Moreover if $q_2 = \infty$,

$$\|u\|_{\infty, R_\infty} \leq 3 \cdot 2^{\frac{q_1 - q}{q}} \left[\left(4^\gamma \frac{q_1}{q} \right)^\gamma \frac{K}{(R_0 - R_\infty)^\gamma} \right]^{\frac{q_1}{q}} \|u\|_{q, R_0}.$$

Proof. See Lemma 3.7 (Extending Local Upper Bounds) of [4]. □

8.3 - Numerical identities and inequalities

Now, in order to be complete and to simplify the reading of this paper, we recall some numerical identities and inequalities used in the paper.

$$(76) \quad \sum_{j=1}^{\infty} s^j = \frac{s}{1 - s}, \quad \forall 0 \leq s < 1 \implies \sum_{j=1}^{\infty} \left(\frac{p}{r}\right)^j = \frac{p}{r - p}, \text{ for } r > p.$$

$$\sum_{j=1}^{\infty} j^N s^j = \left[s \frac{d}{ds} \right]^{(N)} \left(\frac{1}{1 - s} \right), \quad \forall 0 \leq s < 1, \quad N \in \mathbb{N}$$

and so

$$(77) \quad \sum_{j=1}^{\infty} j s^j = \frac{s}{(1 - s)^2}, \quad \forall 0 \leq s < 1 \implies \sum_{j=1}^{\infty} j \left(\frac{p}{r}\right)^j = \frac{pr}{(r - p)^2},$$

for $r > p$.

$$(78) \quad \sum_{j=1}^k s^j = \frac{s(1-s^k)}{1-s}, \quad \forall 0 \leq s < 1 \implies \sum_{j=1}^k \left(\frac{p}{r}\right)^j = \frac{p}{r-p} \left[1 - \left(\frac{p}{r}\right)^k\right],$$

for $r > p$.

$$\sum_{j=k+1}^{\infty} s^j = \frac{s}{1-s} s^k \quad \forall 0 \leq s < 1 \implies \sum_{j=k+1}^{\infty} \left(\frac{p}{r}\right)^j = \frac{p}{r-p} \left(\frac{p}{r}\right)^k,$$

for $r > p$.

Stirling's formula:

$$n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\alpha_n} \quad \text{with} \quad \frac{1}{12n+1} \leq \alpha_n \leq \frac{1}{12n}.$$

ε -version of Young's inequality:

$$(79) \quad a \cdot b \leq \frac{\varepsilon}{\sigma} a^\sigma + \frac{\sigma-1}{\sigma} \frac{b^{\frac{\sigma}{\sigma-1}}}{\varepsilon^{\frac{1}{\sigma-1}}},$$

for any $\varepsilon > 0$, $a, b \geq 0$ and $\sigma > 1$.

Acknowledgments. The first author has been partially funded by Project MTM2011-24696 (Spain). The second author has been supported by the ERC grant 207573 Vectorial Problems. The authors would like to thank the referee for careful reading and suggestions.

References

- [1] F. ANDREU-VAILLO, V. CASELLES and J. M. MAZÓN, *Parabolic quasilinear equations minimizing linear growth functionals*, Birkhäuser Verlag, Basel 2004.
- [2] M. F. BIDAUT-VÉRON, *Local and global behavior of solutions of quasilinear equations of Emden-Fowler type*, Arch. Rational Mech. Anal. **107** (1989), 293-324.
- [3] M. BONFORTE, G. GRILLO and J. L. VÁZQUEZ, *Behaviour near extinction for the Fast Diffusion Equation on bounded domains*, J. Math. Pures Appl. **97** (2012), 1-38.
- [4] M. BONFORTE, G. GRILLO and J. L. VÁZQUEZ, *Quantitative local bounds for subcritical semilinear elliptic equations*, Milan J. Math. **80** (2012), 65-118.
- [5] M. BONFORTE, G. GRILLO and J. L. VÁZQUEZ, *Quantitative bounds for subcritical semilinear elliptic equations*, in "Recent Trends in Nonlinear Partial

- Differential Equations II: Stationary Problems”, Contemp. Math., 595, Amer. Math. Soc., Providence, RI 2013 (in press).
- [6] M. BONFORTE, R. G. IAGAR and J. L. VÁZQUEZ, *Local smoothing effects, positivity, and Harnack inequalities for the fast p -Laplacian equation*, Adv. Math. **224** (2010), 2151-2215.
- [7] M. BONFORTE and J. L. VÁZQUEZ, *Global positivity estimates and Harnack inequalities for the fast diffusion equation*, J. Funct. Anal. **240** (2006), 399-428.
- [8] M. BONFORTE and J. L. VÁZQUEZ, *Reverse smoothing effects, fine asymptotics, and Harnack inequalities for fast diffusion equations*, Bound. Value Probl. **2007**, Art. ID 21425, 31 pp.
- [9] M. BONFORTE and J. L. VÁZQUEZ, *Positivity, local smoothing, and Harnack inequalities for very fast diffusion equations*, Adv. Math. **223** (2010), 529-578.
- [10] H. BRÉZIS and L. NIRENBERG, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), 437-477.
- [11] H. BRÉZIS and R. E. L. TURNER, *On a class of superlinear elliptic problems*, Comm. Partial Differential Equations **2** (1977), 601-614.
- [12] C. BUDD and J. NORBURY, *Semilinear elliptic equations and supercritical growth*, J. Differential Equations **68** (1987), 169-197.
- [13] X. CABRÉ, A. CAPELLA and M. SANCHÓN, *Regularity of radial minimizers of reaction equations involving the p -Laplacian*, Calc. Var. Partial Differential Equations **34** (2009), 475-494.
- [14] X. CABRÉ and M. SANCHÓN, *Semi-stable and extremal solutions of reaction equations involving the p -Laplacian*, Commun. Pure Appl. Anal. **6** (2007), 43-67.
- [15] L. A. CAFFARELLI, B. GIDAS and J. SPRUCK, *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math. **42** (1989), 271-297.
- [16] L. DAMASCELLI, A. FARINA, B. SCIUNZI and E. VALDINOCI, *Liouville results for m -Laplace equations of Lane-Emden-Fowler type*, Ann. Inst. H. Poincaré Anal. Non Linéair **26** (2009), 1099-1119.
- [17] L. DAMASCELLI and B. SCIUNZI, *Regularity, monotonicity and symmetry of positive solutions of m -Laplace equations*, J. Differential Equations **206** (2004), 483-515.
- [18] L. DAMASCELLI and B. SCIUNZI, *Harnack inequalities, maximum and comparison principles, and regularity of positive solutions of m -Laplace equations*, Calc. Var. Partial Differential Equations **25** (2006), 139-159.
- [19] E. N. DANCER, *On the influence of domain shape on the existence of large solutions of some superlinear problems*, Math. Ann. **285** (1989), 647-669.
- [20] E. N. DANCER, *Some notes on the method of moving planes*, Bull. Austral. Math. Soc. **46** (1992), 425-434.
- [21] E. N. DANCER, *Superlinear problems on domains with holes of asymptotic shape and exterior problems*, Math. Z. **229** (1998), 475-491.
- [22] J. DÁVILA, L. DUPAIGNE and A. FARINA, *Partial regularity of finite Morse index solutions to the Lane-Emden equation*, J. Funct. Anal. **261** (2011), 218-232.

- [23] D. G. DE FIGUEIREDO, P.-L. LIONS and R. D. NUSSBAUM, *A priori estimates and existence of positive solutions of semilinear elliptic equations*, J. Math. Pures Appl. **61** (1982), 41-63.
- [24] M. DEL PINO, M. MUSSO and F. PACARD, *Boundary singularities for weak solutions of semilinear elliptic problems*, J. Funct. Anal. **253** (2007), 241-272.
- [25] E. DIBENEDETTO, *$C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations*, Nonlinear Anal. **7** (1983), 827-850.
- [26] E. DIBENEDETTO, U. GIANAZZA and V. VESPRI, *Harnack type estimates and Hölder continuity for non-negative solutions to certain sub-critically singular parabolic partial differential equations*, Manuscripta Math. **131** (2010), 231-245.
- [27] E. DIBENEDETTO, U. GIANAZZA and V. VESPRI, *Harnack's inequality for degenerate and singular parabolic equations*, Springer Monographs in Mathematics, Springer, New York 2012.
- [28] A. FARINA, *Liouville-type results for solutions of $-\Delta u = |u|^{p-1}u$ on unbounded domains of \mathbb{R}^N* , C. R. Math. Acad. Sci. Paris **341** (2005), 415-418.
- [29] A. FARINA, *On the classification of solutions of the Lane-Emden equation on unbounded domains of \mathbb{R}^N* , J. Math. Pures Appl. **87** (2007), 537-561.
- [30] A. FERRERO, *On the solutions of quasilinear elliptic equations with a polynomial-type reaction term*, Adv. Differential Equations **9** (2004), 1201-1234.
- [31] B. GIDAS and J. SPRUCK, *Global and local behavior of positive solutions of nonlinear elliptic equations*, Comm. Pure Appl. Math. **34** (1981), 525-598.
- [32] B. GIDAS and J. SPRUCK, *A priori bounds for positive solutions of nonlinear elliptic equations*, Comm. Partial Differential Equations **6** (1981), 883-901.
- [33] D. GILBARG and N. S. TRUDINGER, *Elliptic partial differential equations of second order*, Grundlehren der Mathematischen Wissenschaften, **224**, Springer-Verlag, Berlin-New York 1977.
- [34] E. GIUSTI, *Direct methods in the calculus of variations*, World Scientific Publishing Co., River Edge, NJ 2003.
- [35] D. D. JOSEPH and T. S. LUNDGREN, *Quasilinear Dirichlet problems driven by positive sources*, Arch. Rational Mech. Anal. **49** (1972/73), 241-269.
- [36] O. A. LADYZHNSKAYA and N. N. URAL'TSEVA, *Linear and quasilinear elliptic equations*, Translated from the Russian by Scripta Technica, Inc. Translation editor: Leon Ehrenpreis, Academic Press, New York-London 1968.
- [37] G. M. LIEBERMAN, *Boundary regularity for solutions of degenerate elliptic equations*, Nonlinear Anal. **12** (1988), 1203-1219.
- [38] C. B. MORREY, *Multiple integrals in the calculus of variations*, Springer-Verlag, New York 1966.
- [39] M. K. V. MURTHY and G. STAMPACCHIA, *Boundary value problems for some degenerate-elliptic operators*, Ann. Mat. Pura Appl. **80** (1968), 1-122.
- [40] R. MAZZEO and F. PACARD, *A construction of singular solutions for a semilinear elliptic equation using asymptotic analysis*, J. Differential Geom. **44** (1996), 331-370.
- [41] L. NIRENBERG, *On elliptic partial differential equations*, Ann. Scuola Norm. Sup. Pisa **13** (1959), 115-162.
- [42] F. PACARD, *Existence de solutions faibles positives de $-\Delta u = u^2$ dans des ouverts bornés de \mathbb{R}^n , $n \geq 3$* , C. R. Acad. Sci. Paris Sér. I Math. **315** (1992), 793-798.

- [43] F. PACARD, *A note on the regularity of weak solutions of $-\Delta u = u^\alpha$ in \mathbf{R}^n , $n \geq 3$* , Houston J. Math. **18** (1992), 621-632.
- [44] F. PACARD, *Existence and convergence of positive weak solutions of $-\Delta u = u^{n/(n-2)}$ in bounded domains of \mathbf{R}^n , $n \geq 3$* , Calc. Var. Partial Differential Equations **1** (1993), 243-265.
- [45] F. PACARD, *Partial regularity for weak solutions of a nonlinear elliptic equation*, Manuscripta Math. **79** (1993), 161-172.
- [46] P. POLÁČIK, P. QUITTNER and P. SOUPLET, *Singularity and decay estimates in superlinear problems via Liouville-type theorems. I. Elliptic equations and systems*, Duke Math. J. **139** (2007), 555-579.
- [47] P. PUCCI and J. SERRIN, *A general variational identity*, Indiana Univ. Math. J. **35** (1986), 681-703.
- [48] P. PUCCI and J. SERRIN, *The strong maximum principle revisited*, J. Differential Equations **196** (2004), 1-66.
- [49] P. PUCCI and J. SERRIN, *The maximum principle*, Progress in Nonlinear Differential Equations and their Applications, **73**, Birkhäuser Verlag, Basel 2007.
- [50] W. RUDIN, *Real and complex analysis*, 2nd ed., McGraw-Hill Book Co., New York-Düsseldorf-Johannesburg 1974.
- [51] J. SERRIN, *Local behavior of solutions of quasi-linear equations*, Acta Math. **111** (1964), 247-302.
- [52] J. SERRIN and H. ZOU, *Cauchy-Liouville and universal boundedness theorems for quasilinear elliptic equations and inequalities*, Acta Math. **189** (2002), 79-142.
- [53] N. S. TRUDINGER, *On Harnack type inequalities and their application to quasilinear elliptic equations*, Comm. Pure Appl. Math. **20** (1967), 721-747.
- [54] N. S. TRUDINGER, *Harnack inequalities for nonuniformly elliptic divergence structure equations*, Invent. Math. **64** (1981), 517-531.
- [55] J. L. VÁZQUEZ, *A strong maximum principle for some quasilinear elliptic equations*, Appl. Math. Optim. **12** (1984), 191-202.

MATTEO BONFORTE

Departamento de Matemáticas
 Universidad Autónoma de Madrid
 Campus de Cantoblanco
 Madrid, 28049, Spain
 e-mail: matteo.bonforte@uam.es

AGNESE DI CASTRO

Dipartimento di Matematica e Informatica
 Università degli Studi di Parma
 Campus - Parco Area delle Scienze, 53/A
 Parma, 43124, Italy
 e-mail: agnese.dicastro@unipr.it
 Dipartimento di Matematica, Università di Pisa
 Largo Bruno Pontecorvo, 5
 Pisa, 56127, Italy
 e-mail: dicastro@mail.dm.unipi.it

